

Ophthalmic Ointment Monographs: Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment

Type of Posting	Revision Bulletin
Posting Date	29-Jul-2016
Official Date	01-Aug-2016
Expert Committee	Chemical Medicines Monographs 1 to 6
Reason for Revision	Compliance

In accordance with the Rules and Procedures of the 2015-2020 Council of Experts, the Chemical Medicines Expert Committees 1 to 6 has revised the monographs listed below. The purpose of the revision is to replace the requirement to comply with the entire content of the USP general chapter *Ophthalmic Products—Quality Tests <771>* with a requirement to comply only with the subsection for *Particulate and Foreign Matter* in *Ophthalmic Products—Quality Tests <771>*, and with the section for *Container Content* for those monographs where the requirement for Minimum Fill was deleted.

- Atropine Sulfate Ophthalmic Ointment
- Bacitracin Ophthalmic Ointment
- Bacitracin Zinc and Polymyxin B Sulfate Ophthalmic Ointment
- Bland Lubricating Ophthalmic Ointment
- Chloramphenicol and Polymyxin B Sulfate Ophthalmic Ointment
- Chloramphenicol Ophthalmic Ointment
- Chlortetracycline Hydrochloride Ophthalmic Ointment
- Ciprofloxacin Ophthalmic Ointment
- Dexamethasone Sodium Phosphate Ophthalmic Ointment
- Erythromycin Ophthalmic Ointment
- Gentamicin and Prednisolone Acetate Ophthalmic Ointment
- Gentamicin Sulfate Ophthalmic Ointment
- Hydrocortisone Acetate Ophthalmic Ointment
- Idoxuridine Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates and Bacitracin Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Ophthalmic Ointment
- Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment
- Neomycin Sulfate and Dexamethasone Sodium Phosphate Ophthalmic Ointment
- Neomycin Sulfate Ophthalmic Ointment
- Oxytetracycline Hydrochloride and Polymyxin B Sulfate Ophthalmic Ointment
- Sodium Chloride Ophthalmic Ointment
- Sulfacetamide Sodium and Prednisolone Acetate Ophthalmic Ointment
- Sulfacetamide Sodium Ophthalmic Ointment
- Tetracycline Hydrochloride Ophthalmic Ointment
- Tobramycin and Dexamethasone Ophthalmic Ointment
- Tobramycin Ophthalmic Ointment

The Revision Bulletins for the monographs listed above supersede the currently official version of these monographs. The Revision Bulletin will be incorporated in the *First Supplement to USP 40–NF 35*.

Should you have any questions, please contact Margareth R. C. Marques, M.Sc., Ph.D. (301-816-8106 or mrm@usp.org).

Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment

DEFINITION

Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment is a sterile ointment containing Neomycin Sulfate, Polymyxin B Sulfate, Bacitracin Zinc, and Hydrocortisone Acetate. It contains the equivalent of NLT 90.0% and NMT 140.0% of the labeled amounts of neomycin, polymyxin B, and bacitracin, and NLT 90.0% and NMT 110.0% of the labeled amount of hydrocortisone acetate (C₂₃H₃₂O₆).

IDENTIFICATION

- **A. THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST** (201BNP): Meets the requirements
- **B.** The retention time of the hydrocortisone acetate peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay for Hydrocortisone Acetate*.

ASSAY

• NEOMYCIN

(See *Antibiotics—Microbial Assays* <81>.)

Sample solution: Shake a portion of Ophthalmic Ointment in a separator with 50 mL of ether. Extract with four 20-mL portions of *Buffer B.3*. Combine the aqueous extracts, and dilute with *Buffer B.3* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.3* to obtain a *Test Dilution* having a neomycin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• POLYMYXIN B

(See *Antibiotics—Microbial Assays* <81>.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 25-mL portions of *Buffer B.6*. Combine the aqueous extracts, and dilute with *Buffer B.6* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.6* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (10 polymyxin B Units/mL). Add to each *Test Dilution* of the standard a quantity of USP Neomycin Sulfate RS, dissolved in *Buffer B.6*, to obtain the same concentration of neomycin as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

• BACITRACIN

(See *Antibiotics—Microbial Assays* <81>.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 25-mL portions of 0.01 N hydrochloric acid. Combine the acid extracts, and dilute with 0.01 N hydrochloric acid to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.1* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (1.0 bacitracin Unit/mL). If the *Sample solution* has a concentration of less than 100 bacitracin Units/mL, add hydrochloric acid to each *Test Dilution* of the standard to obtain the same concentration of hydrochloric acid as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

Change to read:

• HYDROCORTISONE ACETATE

Mobile phase: Butyl chloride, water-saturated butyl chloride, tetrahydrofuran, methanol, and glacial acetic acid (475:475:70:35:30)

Standard solution: 0.10 mg/mL of USP Hydrocortisone Acetate RS in water-saturated chloroform

Sample solution: Nominally 0.10 mg/mL of hydrocortisone acetate from Ophthalmic Ointment prepared as follows. Transfer a portion of Ophthalmic Ointment containing nominally 2.5 mg of hydrocortisone acetate to a closable container. Add 25.0 mL of water-saturated chloroform and about 10 glass beads. Securely close the container, and shake vigorously for approximately 15 min. Centrifuge, and use the clear, lower chloroform layer.

Chromatographic system

(See *Chromatography* <621>, *System Suitability*.)

Mode: LC

Detector: UV 254 nm

Column: ▲3.9-mm × 30-cm; 10-μm▲^{USP39} packing L3

System suitability

Sample: *Standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydrocortisone acetate (C₂₃H₃₂O₆) in the portion of Ophthalmic Ointment taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Hydrocortisone Acetate RS in the *Standard solution* (mg/mL)

C_U = nominal concentration of hydrocortisone acetate in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Delete the following:

- ▲• **MINIMUM FILL** <755>: Meets the requirements

▲^{USP39}

SPECIFIC TESTS

Delete the following:

- ▲• **WATER DETERMINATION, Method I** <921>

Analysis: Use 20 mL of a mixture of toluene and methanol (7:3) in place of methanol in the titration vessel.

Acceptance criteria: NMT 0.5%

▲^{USP39}

Change to read:

- **STERILITY TESTS** <71>: ▲Meets the requirements▲^{USP39}

Delete the following:

- ▲• **METAL PARTICLES IN OPHTHALMIC OINTMENTS** <751>: It meets the requirements.▲^{USP39}

2 Neomycin

Change to read:

▲● **OTHER REQUIREMENTS:** It meets the requirements ● for *Particulate and Foreign Matter and Container Contents* ● (RB 1-Aug-2016) in *Ophthalmic Products—Quality Tests* (771), ● *Drug Product Quality, Universal Tests, Particulate and Foreign Matter and Container Contents*. ● (RB 1-Aug-2016)

▲USP39

ADDITIONAL REQUIREMENTS

Change to read:

- **PACKAGING AND STORAGE:** Preserve in collapsible ophthalmic ointment tubes. ▲Store at controlled room temperature.▲USP39
- **USP REFERENCE STANDARDS** (11)
 - USP Bacitracin Zinc RS
 - USP Hydrocortisone Acetate RS
 - USP Neomycin Sulfate RS
 - USP Polymyxin B Sulfate RS