

## Salix Species Bark Family of monographs

<b>Type of Posting</b>	Notice of Intent to Revise
<b>Posting Date</b>	25–Aug–2017
<b>Official Date</b>	01–Dec–2017, Revision Bulletin and 01–Mar–2018, Interim Revision Announcement
<b>Expert Committee</b>	Botanical Dietary Supplements and Herbal Medicines

In accordance with section 7.04 (c) of the 2015–2020 Rules and Procedures of the Council of Experts, this is to provide notice that the Botanical Dietary Supplements and Herbal Medicines Expert Committee intends to revise the *Salix* Species family of monographs, including *Salix* Species Bark, *Salix* Species Bark Powder, and *Salix* Species Bark Dry Extract.

The *Salix* monographs were initially published for comment in *Pharmacopeial Forum* 42(1) [Jan.–Feb. 2016] and subsequently approved for inclusion in the *Second Supplement to USP 40–NF 35*, which will become official on December 1, 2017. Based on safety concerns the Botanical Dietary Supplements and Herbal Medicines Expert Committee has recommended the addition of the following labeling statement to the *Salix* Species family of monographs which were approved for publication in the *Second Supplement to USP 40–NF 35*: **“Dosage forms prepared with this article should bear the following statement: Not for use in children, women who are pregnant or nursing, or by persons with known sensitivity to aspirin.”**

It is anticipated that proposed Interim Revisions Announcements (IRAs) to the *Salix* species monographs will be published in *Pharmacopeial Forum* 43(5) [Sep.–Oct. 2017] pursuant to section 7.02 of the Rules and Procedures. The comment period for these revisions will end on November 30, 2017. Unless otherwise decided by the Botanical Dietary Supplements and Herbal Medicines Expert Committee based on comments received, the IRAs will be posted online on January 26, 2018 and will become official on March 1, 2018.

In addition, it is also anticipated that the Botanical Dietary Supplements and Herbal Medicines Expert Committee will postpone the official date of the *Salix* Species monographs via Revision Bulletins to be posted July 28, 2017 and become official December 1, 2017. The postponements will remain in effect until the IRAs to be published in *Pharmacopeial Forum* 43(5) become official.

Should you have any questions, please contact Anton Bzhelyansky, Scientific Liaison to the Botanical Dietary Supplements and Herbal Medicines Expert Committee at (301-230-6303 or [anb@usp.org](mailto:anb@usp.org)).

**Change to read:**

**Salix Species Bark Powder**

(This monograph is postponed indefinitely.) (RB 1-Dec-2017)

**DEFINITION**

Salix Species Bark Powder consists of Salix Species Bark reduced to fine or very fine powder. It contains NLT 1.50% of total salicylate derivatives, calculated as salicin (C<sub>13</sub>H<sub>18</sub>O<sub>7</sub>) on the dried basis. Salix Species Bark Powder contains NMT 1.50% of free salicin, calculated on the dried basis.

**IDENTIFICATION**

**A. HPTLC FOR ARTICLES OF BOTANICAL ORIGIN (203)**

**Standard solution A:** 1.50 mg/mL of USP Salicin RS in methanol

**Standard solution B:** 30 mg/mL of USP Salix Species Bark Dry Extract RS in methanol. Sonicate for 10 min, centrifuge, and use the supernatant.

**Sample solution A:** Suspend 1000 mg of Salix Species Bark Powder in 10.0 mL of methanol. Sonicate for 10 min, centrifuge, and use the supernatant.

**Sample solution B:** Combine 5.0 mL of Sample solution A with 1.0 mL of 50-mg/mL anhydrous sodium carbonate. Cap tightly and incubate at 60° for 10 min. Centrifuge and use the supernatant.

**Chromatographic system**

**Adsorbent:** Chromatographic silica gel with an average particle size of 5 µm (HPTLC plate)<sup>1</sup>

**Application volume:** 5.0 µL each of Standard solution A, Standard solution B, and Sample solution A and 6.0 µL of Sample solution B as 8-mm bands

**Relative humidity:** Condition the plate to a relative humidity of 33%.

**Temperature:** Ambient, not to exceed 30°

**Developing solvent system:** Ethyl acetate, methanol, and water (77:13:10)

**Developing distance:** 6 cm

**Derivatization reagent:** Sulfuric acid and methanol (1:9). Slowly add sulfuric acid to ice-cold methanol.

**Analysis**

**Samples:** Standard solution A, Standard solution B, Sample solution A, and Sample solution B

Apply the Samples as bands and dry in air. Develop in a saturated chamber, and dry in a current of air for 5 min. Treat with Derivatization reagent, heat at 100° for 5 min, and examine under white light.

**System suitability:** Under white light, the derivatized chromatogram of Standard solution B displays, in the middle third of the plate, three brown bands: the lower corresponds to the salicin band in Standard solution A; the one above it is due to salicortin; the top band is due to tremulacin. A faint band, which may appear between the salicortin and tremulacin bands, is due to tremuloidin. Two darker brown bands are seen in the lower third of the plate: one proximate to the application line; another, more intense, above it.

**Acceptance criteria:** Under white light, the derivatized chromatogram of Sample solution A shows one or several dark bands due to different salicin esters, whose position and intensity are contingent on the Salix species being used. The salicylate bands of interest are predominantly located in the middle third of the plate, demarcated by the salicin and tremulacin bands of Standard solution B. The salicin band in Sample solution A, corresponding to that in Standard solution A, may be faint or not visible. Additional bands may be seen in

Sample solution A and Sample solution B. In Sample solution B, the bands due to salicin esters are not present, while the salicin band corresponding to that in Standard solution A is the principal band observed. The salicin band in Sample solution A is of lower intensity than the corresponding band in Standard solution A. The salicin band in Sample solution B is of comparable or higher intensity than the corresponding band in Standard solution A.

**B. HPLC**

**Analysis:** Proceed as directed in the test for Salicylates Profile and Limit of Free Salicin.

**Acceptance criteria:** The Sample solution exhibits peaks at retention times corresponding to those of salicin and salicin derivatives in the Standard solutions.

**COMPOSITION**

**CONTENT OF SALICIN**

**Diluent:** Methanol and water (1:1)

**Solution A:** 0.01% trifluoroacetic acid

**Solution B:** Acetonitrile

**Mobile phase:** See Table 1.

Table 1

Time (min)	Solution A (%)	Solution B (%)
0	90	10
10	85	15
30	50	50
32	10	90
35	10	90
37	90	10
45	90	10

**Standard solution A:** 0.50 mg/mL of USP Salicin RS in Diluent

**Sample solution:** Accurately weigh about 1.3 g of Salix Species Bark Powder, transfer to a 200-mL round-bottom flask, and add 40 mL of methanol and 3 mL of 1 N sodium hydroxide. Attach the condenser and heat under reflux for 2 h, with intermittent shaking. Allow to cool, neutralize with 3 mL of 1 N hydrochloric acid, and pass through a paper filter into a 100-mL volumetric flask. Wash the round-bottom flask twice with 5-mL aliquots of methanol and filter into the same 100-mL volumetric flask. Adjust with water to volume, mix well, allow to equilibrate to room temperature, and re-adjust with water. Pass through a PTFE filter of 0.45-µm pore size, discarding the initial 3 mL of the filtrate.

**Chromatographic system**

(See Chromatography (621), System Suitability.)

**Mode:** LC

**Detector:** UV 270 nm

**Column:** 4.6-mm × 25-cm; 5-µm base-deactivated packing L1

**Column temperature:** 30°

**Flow rate:** 1.0 mL/min

**Injection volume:** 10 µL

**System suitability**

**Sample:** Standard solution A

**Suitability requirements**

**Tailing factor:** 0.8–2.0 for the salicin peak

**Relative standard deviation:** NMT 2.0% for the salicin peak in replicate injections

**Analysis**

**Samples:** Standard solution A and Sample solution  
Using the chromatogram of the Standard solution A, identify the salicin peak in the Sample solution chromatogram.

<sup>1</sup> A suitable commercially available plate is HPTLC Silica Gel 60 F<sub>254</sub> from EM D Millipore (e.g., Part No. 1.05642.0001).

## 2 Salix Species

Calculate the percentage of salicin hydrolytically derived from constituent salicylates in the portion of *Salix* Species Bark Powder taken:

$$\text{Result} = (r_u/r_s) \times C_s \times (V/W) \times 100$$

- $r_u$  = peak area of salicin from the *Sample solution*  
 $r_s$  = peak area of salicin from the *Standard solution A*  
 $C_s$  = concentration of USP Salicin RS in the *Standard solution A* (mg/mL)  
 $V$  = volume of the *Sample solution* (mL)  
 $W$  = weight of *Salix* Species Bark Powder taken to prepare the *Sample solution* (mg)

**Acceptance criteria:** NLT 1.50% of salicin on the dried basis

### CONTAMINANTS

- **ARTICLES OF BOTANICAL ORIGIN** (561), *Limits of Elemental Impurities*: Meets the requirements
- **ARTICLES OF BOTANICAL ORIGIN** (561), *Pesticide Residue Analysis*: Meets the requirements
- **MICROBIAL ENUMERATION TESTS** (2021): The total aerobic bacterial count does not exceed  $10^5$  cfu/g, the total combined yeasts and molds count does not exceed  $10^3$  cfu/g, and the bile-tolerant Gram-negative bacteria count does not exceed  $10^3$  cfu/g.
- **ABSENCE OF SPECIFIED MICROORGANISMS** (2022), *Test Procedures, Test for Absence of Salmonella Species* and *Test for Absence of Escherichia coli*: Meets the requirements

### SPECIFIC TESTS

- **SALICYLATES PROFILE AND LIMIT OF FREE SALICIN**<sup>2</sup>  
**Diluent, Mobile phase, Standard solution A, and Chromatographic system:** Proceed as directed in the test for *Content of Salicin*.

**Standard solution B:** 5 mg/mL of USP *Salix* Species Bark Dry Extract RS in *Diluent*. Sonicate for 5 min, mix well, and pass through a PTFE filter of 0.45- $\mu$ m pore size, discarding the initial 3 mL of the filtrate.

**Sample solution:** Accurately weigh about 650 mg of *Salix* Species Bark Powder, transfer to a 50-mL volumetric flask, add 25 mL of methanol, and sonicate for 30 min. Adjust with water to volume, mix well, allow to equilibrate to room temperature, and readjust with water. Pass through a PTFE filter of 0.45- $\mu$ m pore size, discarding the initial 3 mL of the filtrate.

#### System suitability

**Samples:** *Standard solution A* and *Standard solution B*  
**Suitability requirements**

**Tailing factor:** 0.8–2.0 for the salicin peak, *Standard solution A*

**Relative standard deviation:** NMT 2.0% determined for the salicin peak in replicate injections, *Standard solution A*

**Chromatographic similarity:** The chromatogram is similar to the reference chromatogram provided with the lot of USP *Salix* Species Bark Dry Extract RS being used, *Standard solution B*.

#### Analysis

**Samples:** *Standard solution A*, *Standard solution B*, and *Sample solution*

Using the chromatograms of *Standard solution A*, *Standard solution B*, and the reference chromatogram provided with the lot of USP *Salix* Species Bark Dry Extract RS being used, identify salicin esters present in the *Sample solution* chromatogram. The approximate relative retention times, with respect to salicin, are provided in *Table 2*.

**Table 2**

Analyte	Relative Retention Time
Salicin	1.0
Salicortin	3.0
Tremuloidin	3.6
Tremulacin	4.6

Calculate the percentage of salicin in the portion of *Salix* Species Bark Powder taken:

$$\text{Result} = (r_u/r_s) \times C_s \times (V/W) \times 100$$

- $r_u$  = peak area of salicin from the *Sample solution*  
 $r_s$  = peak area of salicin from *Standard solution A*  
 $C_s$  = concentration of USP Salicin RS in *Standard solution A* (mg/mL)  
 $V$  = volume of the *Sample solution* (mL)  
 $W$  = weight of *Salix* Species Bark Powder taken to prepare the *Sample solution* (mg)

**Acceptance criteria:** NMT 1.50% of salicin on the dried basis. The peak area of salicin is NMT 50% of the combined peak areas of all identified constituent salicylates.

### BOTANICAL CHARACTERISTICS

**Macroscopic:** Pale yellow, greenish-yellow, or light brown powder

**Microscopic:** Bundles of narrow fibers, up to about 600  $\mu$ m long, with very thick walls, lignified, and surrounded by a crystal sheath containing prism crystals of calcium oxalate; parenchyma of the cortex with thick, pitted and deeply beaded walls, and containing large cluster crystals of calcium oxalate; uniseriate medullary rays; thickened and suberized cork cells. Groups of brownish collenchyma from the bud may be present. Twigs show fragments of lignified fibers and vessels from the xylem.

### LOSS ON DRYING (731)

**Sample:** 1.0 g of *Salix* Species Bark Powder

**Analysis:** Dry the *Sample* at 105° for 2 h.

**Acceptance criteria:** NMT 10.0%

### ARTICLES OF BOTANICAL ORIGIN (561), Methods of Analysis, Total Ash

**Sample:** 2.0 g of *Salix* Species Bark Powder

**Acceptance criteria:** NMT 10.0%

### ARTICLES OF BOTANICAL ORIGIN (561), Methods of Analysis, Acid-Insoluble Ash

**Sample:** 2.0 g of *Salix* Species Bark Powder

**Acceptance criteria:** NMT 3.0%

### ARTICLES OF BOTANICAL ORIGIN (561), Methods of Analysis, Water-Soluble Extractives

**Sample:** 2.0 g of *Salix* Species Bark Powder

**Acceptance criteria:** NLT 10.0%

### ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers, protected from light and moisture, and store at room temperature.
- **LABELING:** The label states the Latin binomial(s) of one or several *Salix* species included in the article.
- **USP REFERENCE STANDARDS** (11)

USP Salicin RS

USP *Salix* Species Bark Dry Extract RS

• (This monograph is postponed indefinitely.) • (RB 1-Dec-2017)

■2S (USP40)

<sup>2</sup>Elevated free salicin content may indicate prehydrolysis or fortification with extraneous salicin.