

[x](#)

How to Use

- **Searching:** Type keyword in search field at top of page. Search by all or part of a monograph title. For searches using multiple criteria, you will find items that match each of the specified criteria unless quotation marks are used.
 - For example, a search on Aminosalicyclic Acid Tablets will result in anything that contains “Aminosalicyclic” OR “Acid” OR “Tablets”
 - A search for “Aminosalicyclic Acid Tablets” will result in anything that specifically contains “Aminosalicyclic Acid Tablets”
- **Sorting:** Click on any column header title to sort alphabetically or chronologically in ascending or descending order. Note: the page load column is sorted alphabetically so that a number is ordered by first digit vs. by the actual number; thus, numbers will not always be in order.
 - For example, page 2178 will come before page 74 on a page sort.
- **Downloading:** You can download the Errata table in Comma-separated Value (.csv). The download will include the Errata that you have filtered on.
- **Importing:** You will need to import the file into Excel or Open Office with UTF-8 encoding, as opposed to simply opening it. To import, open Excel or Open Office and select import from the File drop-down. Depending on the version you are using, you should be presented with import formatting options to include UTF-8 as one of the first steps. Importing via UTF-8 should eliminate odd character conversions.

Monograph Title	Section	Source	Page Number	Errata Post	Errata Official	Target Errata	Target Online	Description
		Publication		Date	Date	Print Publication	Fix Publication	
DIDANOSINE	ASSAY/	USP41–NF36	1275	27-Apr-2018	1-May-2018	USP42–NF37	Second	Line 1 of <i>Guard</i>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
FOR ORAL SOLUTION	<i>Procedure/Chromatographic system</i>							<i>Supplement to USP41–NF36</i>	<i>column: Change 20-cm; to: 20-mm;</i>
ISOSORBIDE DINITRATE SUBLINGUAL TABLETS	<i>Assay</i>	<i>USP41–NF36</i>	<i>2272</i>	<i>27-Apr-2018</i>		<i>1-May-2018</i>	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	<i>Change Buffer solution, Mobile phase, Internal standard solution, Standard preparation, and Chromatographic system—Prepare as directed in the Assay under Diluted Isosorbide Dinitrate. to: Buffer solution —Dissolve 15.4 g of ammonium acetate in water, add 11.5 mL of glacial acetic acid,</i>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>dilute with water to 1000 mL, and mix to obtain a solution having a pH of about 4.7.</p> <p><i>Mobile phase</i>—Mix 350 mL of water, 100 mL of <i>Buffer solution</i>, and 550 mL of methanol. Cool to room temperature, dilute with water to 1000 mL, mix, degas, and filter. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> <621>).</p> <p><i>Internal standard solution</i> —Transfer a quantity of diluted</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>nitroglycerin to a suitable volumetric flask, add about 60% of the flask volume of methanol, sonicate for 5 minutes, and shake for 30 minutes. Dilute with methanol to volume to obtain a solution having a concentration of about 3 mg of nitroglycerin per mL, and mix. Allow any undissolved material to settle, filter, and store the filtrate in an airtight container.</p> <p><i>Standard preparation</i></p> <p>—Transfer about 125 mg of</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							recently mixed USP Diluted Isosorbide Dinitrate RS, accurately weighed, to a 50-mL volumetric flask, add about 30 mL of <i>Mobile phase</i> , shake for 30 minutes, dilute with <i>Mobile phase</i> to volume, and mix. Pipet 10 mL of the resulting solution into a 25-mL volumetric flask, and add 4.0 mL of <i>Internal standard solution</i> and 4 mL of dilute <i>Buffer solution</i> (1 in 10). Cool to room temperature, dilute with

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p><i>Mobile phase to volume, and mix to obtain a solution having a known concentration of about 0.25 mg of isosorbide dinitrate per mL, based on the quantity of USP Diluted Isosorbide Dinitrate RS weighed and the labeled content of isosorbide dinitrate. Pass a portion of this solution through a 0.45-μm filter.</i></p> <p>AND</p> <p>Add</p> <p><i>Chromatographic system (see Chromatography <621>)</i>—The liquid chromatograph is equipped with</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>a 220-nm detector and a 4-mm x 25-cm column that contains packing L1. The flow rate is about 1 mL per minute.</p> <p>Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the resolution, <i>R</i>, between isosorbide dinitrate and nitroglycerin is not less than 2.0; and the relative standard deviation for replicate injections determined from the peak response ratios</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>is not more than 2%. [Note—The relative retention times are about 0.75 for isosorbide dinitrate and 1.0 for nitroglycerin. The relative retention times for isosorbide mononitrates, if present, are about 0.38.]</p> <p>AND</p> <p>Line 1 of <i>Procedure:</i> Change Proceed as directed for <i>Procedure</i> in the Assay under <i>Diluted Isosorbide Dinitrate</i>. to: Separately inject equal volumes (about 20 µL) of the <i>Standard</i></p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
									<p><i>preparation and the Assay preparation into the chromatograph, record the chromatograms, and measure the responses for the major peaks.</i></p> <p>Line 1 of <i>0.1 N Potassium Hydroxide VS: Change Transfer 100 mL of potassium hydroxide to a 1000-mL volumetric flask. to: Transfer 100 mL of 1 N Potassium Hydroxide VS to a 1000-mL volumetric flask.</i></p>
REAGENTS	SOLUTIONS/ Volumetric Solutions	USP41–NF36	5769	27-Apr-2018		1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	
REPOSITORY	ADDITIONAL REQUIREMENTS	USP41–NF36	1097	27-Apr-2018		1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	In USP Reference Standards

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ISOSORBIDE Assay DINITRATE CHEWABLE TABLETS	USP41–NF36	2270	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	<11>: Add USP Ascorbic Acid RS Change Buffer solution, Mobile phase, Internal standard solution, Standard preparation, and Chromatographic system—Prepare as directed in the Assay under Diluted Isosorbide Dinitrate. to: Buffer solution —Dissolve 15.4 g of ammonium acetate in water, add 11.5 mL of glacial acetic acid, dilute with water to 1000 mL, and

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>mix to obtain a solution having a pH of about 4.7.</p> <p><i>Mobile phase</i>—Mix 350 mL of water, 100 mL of <i>Buffer solution</i>, and 550 mL of methanol. Cool to room temperature, dilute with water to 1000 mL, mix, degas, and filter. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> <621>.)</p> <p><i>Internal standard solution</i> —Transfer a quantity of diluted nitroglycerin to a suitable</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>volumetric flask, add about 60% of the flask volume of methanol, sonicate for 5 minutes, and shake for 30 minutes. Dilute with methanol to volume to obtain a solution having a concentration of about 3 mg of nitroglycerin per mL, and mix. Allow any undissolved material to settle, filter, and store the filtrate in an airtight container.</p> <p><i>Standard preparation</i></p> <p>—Transfer about 125 mg of recently mixed USP Diluted</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>Isosorbide Dinitrate RS, accurately weighed, to a 50-mL volumetric flask, add about 30 mL of <i>Mobile phase</i>, shake for 30 minutes, dilute with <i>Mobile phase</i> to volume, and mix. Pipet 10 mL of the resulting solution into a 25-mL volumetric flask, and add 4.0 mL of <i>Internal standard solution</i> and 4 mL of dilute <i>Buffer solution</i> (1 in 10). Cool to room temperature, dilute with <i>Mobile phase</i> to volume, and</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>mix to obtain a solution having a known concentration of about 0.25 mg of isosorbide dinitrate per mL, based on the quantity of USP Diluted Isosorbide Dinitrate RS weighed and the labeled content of isosorbide dinitrate. Pass a portion of this solution through a 0.45-μm filter.</p> <p>AND</p> <p>Add</p> <p><i>Chromatographic system (see Chromatography <621>)</i>—The liquid chromatograph is equipped with a 220-nm detector and a</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>4-mm x 25-cm column that contains packing L1. The flow rate is about 1 mL per minute.</p> <p>Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the resolution, <i>R</i>, between isosorbide dinitrate and nitroglycerin is not less than 2.0; and the relative standard deviation for replicate injections determined from the peak response ratios is not more than 2%. [Note—The</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>relative retention times are about 0.75 for isosorbide dinitrate and 1.0 for nitroglycerin. The relative retention times for isosorbide mononitrates, if present, are about 0.38.]</p> <p>AND</p> <p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the <i>Assay</i> under <i>Diluted Isosorbide Dinitrate</i>. to: Separately inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the <i>Assay</i></p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Errata Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
									<i>preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks.
REAGENTS	REAGENT SPECIFICATIONS	USP41–NF36	5680	27-Apr-2018		1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 1 of Carbon Disulfide, CS: Change Carbon Disulfide, CS to: Carbon Disulfide, CS ₂
COLCHICINE	IM PURITIES/Limit of Ethyl Acetate/System suitability/Suitability requirements	First Supplement to USP41–NF36	8314	27-Apr-2018		1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Delete Tailing factor. NMT 2.0 for the menthol peak
ARGATROBAN	IM PURITIES/Organic Impurities	USP41–NF36	346	27-Apr-2018		1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Footnote b of Table 2: Change Ethyl (2R,4R)-1-[N ⁸ -nitro-L-arginyl]-

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ISOSORBIDE Assay DINITRATE TABLETS	USP41–NF36	2270	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	4-methylpiperidine-2-carboxylate hydrochloride. to: Ethyl (4R)-1-[N ⁸ -nitro-L-arginyl]-4-methylpiperidine-2-carboxylate. Change Buffer solution, Mobile phase, Internal standard solution, Standard preparation, and Chromatographic system—Prepare as directed in the Assay under Diluted Isosorbide Dinitrate. to: Buffer solution—Dissolve 15.4 g of ammonium

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>acetate in water, add 11.5 mL of glacial acetic acid, dilute with water to 1000 mL, and mix to obtain a solution having a pH of about 4.7.</p> <p><i>Mobile phase</i>—Mix 350 mL of water, 100 mL of <i>Buffer solution</i>, and 550 mL of methanol. Cool to room temperature, dilute with water to 1000 mL, mix, degas, and filter. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> <621>).</p> <p><i>Internal standard</i></p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p><i>solution</i> —Transfer a quantity of diluted nitroglycerin to a suitable volumetric flask, add about 60% of the flask volume of methanol, sonicate for 5 minutes, and shake for 30 minutes. Dilute with methanol to volume to obtain a solution having a concentration of about 3 mg of nitroglycerin per mL, and mix. Allow any undissolved material to settle, filter, and store the filtrate in an airtight container.</p> <p><i>Standard</i></p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p><i>prep aration</i> —Transfer about 125 mg of recently mixed USP Diluted Isosorbide Dinitrate RS, accurately weighed, to a 50-mL volumetric flask, add about 30 mL of <i>Mobile phase</i>, shake for 30 minutes, dilute with <i>Mobile phase</i> to volume, and mix. Pipet 10 mL of the resulting solution into a 25-mL volumetric flask, and add 4.0 mL of <i>Internal standard solution</i> and 4 mL of dilute <i>Buffer solution</i></p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							(1 in 10). Cool to room temperature, dilute with <i>Mobile phase</i> to volume, and mix to obtain a solution having a known concentration of about 0.25 mg of isosorbide dinitrate per mL, based on the quantity of USP Diluted Isosorbide Dinitrate RS weighed and the labeled content of isosorbide dinitrate. Pass a portion of this solution through a 0.45- μ m filter. AND Add <i>Chromatographic system</i> (see <i>Chromatograph</i>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>y <621>—The liquid chromatograph is equipped with a 220-nm detector and a 4-mm × 25-cm column that contains packing L1. The flow rate is about 1 mL per minute. Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the resolution, <i>R</i>, between isosorbide dinitrate and nitroglycerin is not less than 2.0; and the relative standard deviation for replicate</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>injections determined from the peak response ratios is not more than 2%. [Note—The relative retention times are about 0.75 for isosorbide dinitrate and 1.0 for nitroglycerin. The relative retention times for isosorbide mononitrates, if present, are about 0.38.]</p> <p>AND</p> <p>Line 1 of <i>Procedure:</i> Change Proceed as directed for <i>Procedure</i> in the Assay under <i>Diluted Isosorbide Dinitrate.</i> to: Separately</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ZONISAMIDE CAPSULES	PERFORMANCE TESTS/ <i>Dissolution</i> <711>	USP41–NF36	4410	27-Apr-2018		1-May-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the <i>Assay preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks. Line 1 of <i>Apparatus 2: Change</i> 75 rpm, with sinkers (see <i>Dissolution</i> <711>, <i>Figure 2a</i>) to: 75 rpm. Use suitable sinkers, if necessary.
REAGENTS/INDICATORS AND SOLUTIONS	SOLUT ION <i>S/Volumetric</i>	USP41–NF36	5772	27-Apr-2018		1-May-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	In the equation in <i>Standardization: Change</i>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							N = mg K ₂ Cr ₂ O ₇ /49.04 x mL Na ₂ S ₂ O ₃ to: M = mg K ₂ Cr ₂ O ₇ /49.04 x mL Na ₂ S ₂ O ₃
DES Loratadine Impurities NE TABLETS	USP41–NF36	1178	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Organic Impurities: Add Protect all solutions containing desloratadine from light.
ISOSORBIDE Dinitrate SUBLINGUAL TABLETS	USP41–NF36	2272	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 1: Change Tablets respond to the Identification test under Isosorbide Dinitrate Tablets. to: Transfer a suitable quantity of finely powdered Tablets to a glass-stoppered centrifuge tube. Add 10 mL of

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>sodium hydroxide solution (1 in 250), shake to wet the powder, then add 15 mL of solvent hexane, and again shake. Centrifuge the mixture, and transfer the upper phase to a beaker. Evaporate the solvent, and dry the residue in vacuum over anhydrous calcium chloride at room temperature for 16 hours: the IR absorption spectrum of a suitable solution in chloroform of the residue so obtained exhibits maxima only at the</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
HYDROGENATED VEGETABLE OIL DEFINITION	USP41–NF36	5649	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	same wavelengths as that of a similar preparation from the residue obtained from USP Diluted Isosorbide Dinitrate RS. Line 2: Change The melting range, heavy metals limit, iodine value, and saponification value differ, to: The melting range, iodine value, and saponification value differ,
PEMETREXED ASSAY/ FOR INJECTION	Procedure/Analysis	Online	27-Apr-2018	1-May-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 8 of the variable definition list: Change M_{r2} = molecular weight of pemetrexed disodium

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
SALMETEROL INHALATION POWDER	ADDITIONAL REQUIREMENT S/USP Reference Standards <11>	USP40–NF35	6096	30-Mar-2018		1-Apr-2018	USP42–NF37	USP42–NF37	(anhydrous), 473.37 to: M_r = molecular weight of pemetrexed disodium (anhydrous), 471.38 Line 2 of USP Salmeterol Related Compound H RS: Change 1-Hydroxy-4-[2-hydroxy-5-(1-hydroxy-2-[[6-(4-p henylbutoxy)hexyl]amino)ethyl]benzyl]-2-napht hoic acid. $C_{36}H_{43}NO_6$ 585.73 to: 1-Hydroxy-4-[2-hydroxy-5-(1-hydroxy-2-[[6-(4-p henylbutoxy)hexyl]amino)ethyl]benzyl]-2-napht hoic acid,

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
FLUTICASONE ADDITIONAL R PROPRIONATE EQUIREMENT AND SALMETEROL INHALATION POWDER		<i>USP40–NF35</i>	4309	30-Mar-2018		1-Apr-2018	<i>USP42–NF37</i>	<i>USP42–NF37</i>	monohydrate. C ₃₆ H ₄₃ NO ₆ · H ₂ O 603.76 Line 2 of USP Salmeterol Related Compound H RS: Change 1-Hydroxy-4-[2- hydroxy-5-(1-hy droxy-2-{{[6-(4-p henylbutoxy)he xyl]amino}ethyl) benzyl]-2-napht hoic acid. C ₃₆ H ₄₃ NO ₆ 585.73 to: 1-Hydroxy-4-[2- hydroxy-5-(1-hy droxy-2-{{[6-(4-p henylbutoxy)he xyl]amino}ethyl) benzyl]-2-napht hoic acid, monohydrate. C ₃₆ H ₄₃ NO ₆ · H ₂ O 603.76
PLASTIC MATERIALS OF CONSTRU	TEST METHODS/ <i>Tin</i> <i>in Non-Tin-</i>	<i>USP41–NF36</i>	6403	30-Mar-2018		1-Apr-2018	<i>USP42–NF37</i>	<i>USP42–NF37</i>	Line 3 of <i>Sample solution:</i>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
CTION	<i>Stabilized Materials</i>								Change If the solution is not colorless, add the sodium sulfate to: If the solution is not colorless, add the sodium sulfite
PERINDOPRIL ERBUMINE	IM PURITIES/ <i>Limit of Perindopril Related Compound I/System suitability</i>	USP40–NF35	5644	30-Mar-2018		1-Apr-2018	USP42–NF37	USP42–NF37	Line 3: Add [Note—The relative retention times for perindopril and perindopril related compound I are 1.0 and 1.6, respectively.]
CHLORHEXIDI NE GLUCONATE ORAL RINSE	<i>Identification</i>	USP40–NF35	3367	23-Feb-2018		1-Mar-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	Line 1 of C: Change Undiluted Oral Rinse used as the test solution meets the requirements for <i>Identification test B</i> under <i>Calcium Gluconate</i> ,

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>except that a Standard solution containing about 0.6 mg of USP Potassium Gluconate RS per mL is used and 15 ?L of the test solution and the Standard solution are applied to the thin-layer chromatographic plate.</p> <p>to:</p> <p>Use undiluted Oral Rinse as the test solution and prepare a Standard solution of USP Potassium Gluconate RS in water containing 0.6 mg/mL. Apply separate 15-?L portions of the</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							test solution and the Standard solution to a suitable thin-layer chromatographic plate (see <i>Chromatography</i> <621>) coated with a 0.25-mm layer of chromatographic silica gel, and allow to dry. Develop the chromatogram in a solvent system consisting of a mixture of alcohol, water, ammonium hydroxide, and ethyl acetate (50:30:10:10) until the solvent front has moved about three-fourths of the length of the

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>plate. Remove the plate from the chamber, and dry at 110° for 20 minutes. Allow to cool and spray with a spray reagent prepared as follows.</p> <p>Dissolve 2.5 g of ammonium molybdate in about 50 mL of 2 N sulfuric acid in a 100-mL volumetric flask, add 1.0 g of ceric sulfate, swirl to dissolve, dilute with 2 N sulfuric acid to volume, and mix. Heat the plate at 110° for about 10 minutes: the principal spot obtained from the test solution corresponds in</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
REAGENTS	REAGENT SPECIFICATIONS	USP41–NF36	5724	23-Feb-2018		1-Mar-2018	USP42–NF37	Second Supplement to USP41–NF36	color, size, and R_F value to that obtained from the Standard solution. Line 2 of 9Z- <i>Retinoic Acid</i> : Change <i>Acidalitretinoin</i>), to: <i>Alitretinoin</i>),
AMLODIPINE AND ATORVASTATIN TABLETS	IM PURITIES/ <i>Organic Impurities Related to Atorvastatin</i>	First Supplement to USP41–NF36	8270	23-Feb-2018		1-Mar-2018	USP42–NF37	Second Supplement to USP41–NF36	In the variable definition list in <i>Analysis</i> : Change M_{r2} = molecular weight of atorvastatin calcium, 1209.39 to: M_{r2} = molecular weight of atorvastatin calcium, 1155.34
PIPERACILLIN AND TAZOBACTAM FOR INJECTION	IM PURITIES/ <i>Organic Impurities, Procedure</i>	USP40–NF35	5728	23-Feb-2018		1-Mar-2018	USP42–NF37	Second Supplement to USP41–NF36	Footnote m: Change (2 <i>S</i> ,5 <i>R</i> ,6 <i>R</i>)-Ethyl 6-((<i>R</i>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>)-2-((2<i>S</i>,5<i>R</i>,6<i>R</i>)-6-[(<i>R</i>)-2-(4-ethyl-2,3-dioxopiperazine-1-carboxamido)-2-phenylacetamido]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxamido)-2-phenylacetamido)-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate.</p> <p>to:</p> <p>2-(((3-Acetyl-4-(ethoxycarbonyl)-5,5-dimethylthiazolidin-2-yl)methyl)amino)-2-oxo-1-phenylethyl 6-(2-(4-ethyl-2,3-dioxopiperazine-1-carboxamido)-2-phenylacetamido)-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[</p>

2/ Table 3

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
AMLODIPINE ASSAY/ AND ATORVAS Procedure TATIN TABLETS	<i>First Supplement to USP41–NF36</i>	8270	23-Feb-2018	1-Mar-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	3.2.0]heptane-2-carboxylate. In the variable definition list of the second equation in <i>Analysis</i> : Change M_{r2} = molecular weight of atorvastatin calcium, 1209.39 to: M_{r2} = molecular weight of atorvastatin calcium, 1155.34
NOREPINEPH IDENTIFICATIO RINE N/B. Procedure BITARTRATE	<i>USP40–NF35</i>	5380	23-Feb-2018	1-Mar-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	Line 1 of <i>Analysis</i> : Change Add 1 drop of ferric chloride TS. to: Add 1 drop of ferric chloride TS to 2 mL of <i>Sample solution</i> .

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
INHALATION AND NASAL DRUG PRODUCTS: AEROSOLS, SPRAYS, AND POWDERS—PERFORMANCE QUALITY TESTS	C. AERODYNAMIC SIZE DISTRIBUTION—INHALATION AEROSOLS, SPRAYS, AND POWDERS	USP41–NF36	6327	23-Feb-2018		1-Mar-2018	USP42–NF37	Second Supplement to USP41–NF36	Figure 6: Change Boquilla del Inhalador to: Inhaler Mouthpiece AND Change Tubo de Admisión to: Induction Port AND Change Cono de Ingreso to: Entrance Cone
ANTIBIOTICS—MICROBIAL ASSAYS	CALCULATION S/Turbidimetric Assay/Sample Data	USP40–NF35	143	23-Feb-2018		1-Mar-2018	USP42–NF37	Second Supplement to USP41–NF36	Second equation in Step 1: Change 0.0125 = to: 0.0062 = AND Third equation in Step 1: Change 0.0325 = to:

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
PAROXETINE HYDROCHLORIDE	ADDITIONAL REQUIREMENT S/USP Reference Standards <11>	<i>First Supplement to USP40–NF35</i>	Online	23-Feb-2018	1-Mar-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	0.0322 = Line 1 of USP Paroxetine Related Compound F RS: Change <i>trans</i> (?) -1-Methyl-3-[1,3-benzodioxol-5-yloxy)methyl]-4-(fluorophenyl)piperidine. to: (3 <i>S</i> ,4 <i>R</i>) -3-[(Benzodioxol-5-yloxy)methyl]-4-(4-fluorophenyl)-1-methylpiperidine.
AMLODIPINE AND ATORVASE TATIN TABLETS	PERFORMANCE TESTS/ Dissolution <711>	<i>First Supplement to USP41–NF36</i>	8270	23-Feb-2018	1-Mar-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	In the variable definition list of the second equation in <i>Analysis</i> : Change M_{r2} = molecular weight of atorvastatin calcium, 1209.39 to:

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
NOREPINEPHRINE BITARTRATE	ASSAY/ <i>Procedure</i>	USP40–NF35	5380	23-Feb-2018		1-Mar-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	<p>M_{r2} = molecular weight of atorvastatin calcium, 1155.34</p> <p>Line 1 of <i>Sample solution</i>: Change 25 mg/mL of Norepinephrine Bitartrate in glacial acetic acid. If necessary warm slightly to effect solution.</p> <p>to: Dissolve 500 mg of Norepinephrine Bitartrate in 20 mL of glacial acetic acid, warming slightly if necessary to effect solution.</p>
STATISTICAL TOOLS FOR PROCEDURE VALIDATION	3. ACCURACY AND PRECISION/3.1 <i>Methods for</i>	USP41–NF36	7622	23-Feb-2018		1-Mar-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	<p>Paragraph 4: Change For example, with $\alpha = 0.05$</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
									and $n = 9$, $t_{0.95:8} = 1.860$ provides a 100(1 - 2 × 1.05)% to: For example, with $\alpha = 0.05$ and $n = 9$, $t_{0.95:8} = 1.860$ provides a 100(1 - 2 × 0.05)%
POWDERED ECHINACEA PALLIDA EXTRACT	IDENTIFICATION	USP40–NF35	6935	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 5 of A. <i>Thin-Layer Chromatography/Presence of echinacoside and absence of dicaffeoylquinic acid/System suitability.</i> Change <i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated,</p> <p>to:</p> <p><i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated,</p> <p>AND</p> <p>Lines 3 and 6 of C.: Change <i>Standard solution B</i>, to:</p> <p><i>Standard solution C</i>,</p> <p>AND</p> <p>Change</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
PHENYTOIN ORAL SUSPENSION	PERFORMANCE TESTS/ <i>Dissolution</i> <711>	USP41–NF36	3286	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Standard solution C) to: Standard solution D) In the Analysis: Change $C_S =$ concentration of USP Phenytoin RS in the Standard solution to: $C_S =$ concentration of USP Phenytoin RS in the Standard solution (mg/mL)
ESZOPICLONE TABLETS	ADDITIONAL REQUIREMENT S/USP Reference Standards <11>	Revision Bulletin (Official August 01, 2017)	Online	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 2 of USP Eszopiclone Related Compound A RS: Change 6-(5-Chloropyridin-2-yl)-7-oxo-6,7-di-hydro-5H-pyr

Monograph TitleSection	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>rolo[3,4 -b]pyrazin-5-yl 4 -methylpiperazi ne-1-carboxylat e 4-oxide. C₁₇H₁₇ClN₆O₄ 404.81 to: [Note—This material may be available in the free base or salt form.] 6-(5-Chloropyrid in-2-yl)-7-oxo-6, 7-di hydro-5<i>H</i> -pyr rolo[3,4 -b]pyrazin-5-yl 4 -methylpiperazi ne-1-carboxylat e 4-oxide. C₁₇H₁₇ClN₆O₄ 404.81 6-(5-Chloropyrid in-2-yl)-7-oxo-6, 7-di hydro-5<i>H</i> -pyr rolo[3,4</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
POWDERED DIGITALIS	IDENTIFICATION N/B. <i>Thin-Layer Chromatographic Identification Test</i>	USP40–NF35	3762	26-Jan-2018		1-Feb-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	<p>-b]pyrazin-5-yl 4-methylpiperazine-1-carboxylate 4-oxide, 3-chlorobenzoic salt (1:1). $C_{17}H_{17}ClN_6O_4 \cdot C_7H_5ClO$ 561.38</p> <p>Line 3 of <i>Standard solution A</i>: Change lead acetate, to: lead acetate TS, AND Line 11 of <i>Analysis</i>: Change Locate the two prominent bands from <i>Standard solution A</i> corresponding in R_F value to the two bands from <i>Standard solution B</i>. to:</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
NEVIRAPINE TABLETS	IMPURITIES/ <i>Organic Impurities</i>	USP40–NF35	5333	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Locate the prominent bands from <i>Standard solution A</i> corresponding in R_f value to the band from <i>Standard solution B</i> . Line 1 of <i>Standard solution</i> : Change 0.125 ?g/mL of USP Nevirapine Anhydrous RS from <i>Standard stock solution A</i> in <i>Diluent</i> to: 0.125 ?g/mL of USP Nevirapine Anhydrous RS in <i>Diluent</i>
ECHINACEA PALLIDA	IDENTIFICATION	USP40–NF35	6931	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 5 of A. <i>Thin-Layer Chromatography/Presence of echinacoside and absence of</i>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p><i>dcaffeoylquinic acid/System suitability.</i></p> <p>Change <i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to:</p> <p><i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
POWDERED ECHINACEA PURPUREA	IDENTIFICATION	USP40–NF35	6942	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	<p>F) that are clearly separated, AND Lines 3 and 6 of C.: Change <i>Standard solution B</i>, to: <i>Standard solution C</i>, AND Change <i>Standard solution C</i>) to: <i>Standard solution D</i>) Line 3 of A. <i>Thin-Layer Chromatography/ Presence of chicoric acid and absence of echinacoside/System suitability</i>. Change <i>Standard</i></p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ECHINACEA SPECIES POWDER	IDENTIFICATIO N	USP41–NF36	4595	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	<p><i>solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to:</p> <p><i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated,</p> <p>Line 4 of A. HPTLC for Articles of</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
CAPSULES							<p><i>Botanical Origin</i> <203>/For Capsules containing <i>Echinacea</i> <i>angustifolia</i> powder prepared from dried rhizome and roots/System suitability: Change <i>Standard</i> <i>solution B</i> shows two major blue bands at about the middle section due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to: <i>Standard</i> <i>solution B</i> shows two major blue</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
TIMOLOL MALEATE	ADDITIONAL R EQUIREMENT S/USP Reference Standards <11>	<i>First Supplement to USP40–NF35</i>	8416	26-Jan-2018		1-Feb-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	bands at about the middle section due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated, Line 2 of USP Timolol Related Compound A RS: Change (R)-1-(<i>tert</i> -Butylamino)-3-(4-morpholino-1,2,5-thiadiazol-3-yloxy)propan-2-ol. $C_{13}H_{24}N_4O_3S$ 316.42 to: (R)-1-(<i>tert</i> -Butylamino)-3-(4-morpholino-1,2,5-thiadiazol-3-yloxy)propan-2-ol maleate. $C_{13}H_{24}N_4O_3S \cdot C$

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p>4H₄O₄ 432.49 AND Line 2 of USP Timolol Related Compound C RS: Change <i>N-(tert</i> -Butyl)-2,3-bis(4 -morpholino-1,2 ,5-thiadiazol-3-y loxy)propan-1-a mine. C₁₉H₃₁N₇O₄S₂ 485.19 to: <i>N-(tert</i> -Butyl)-2,3-bis(4 -morpholino-1,2 ,5-thiadiazol-3-y loxy)propan-1-a mine maleate. C₁₉H₃₁N₇O₄S₂ ? C₄H₄O₄ 601.69 AND Line 3 of USP Timolol Related Compound D RS: Change C₆H₉N₇O₄S to: C₆H₉N₃O₂S</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
IMMUNOLOGICAL TEST PROCEDURES—ENZYMES	USP40–NF35	1344	26-Jan-2018	1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	<p>AND</p> <p>Line 2 of USP Timolol Related Compound E RS: Change (S)-3-(<i>tert</i>-Butylamino)-1-(4-morpholino-1,2,5-thiadiazol-3-yl)oxy]propan-2-yl hydrogen maleate.</p> <p>$C_{17}H_{26}N_4O_6S$ 414.48</p> <p>to:</p> <p>(S,Z)-4-({1-(<i>tert</i>-Butylamino)-3-[(4-morpholino-1,2,5-thiadiazol-3-yl)oxy]propan-2-yl}oxy)-4-oxobut-2-enoic acid maleate salt (1:1)</p> <p>$C_{17}H_{26}N_4O_6S$? $C_4H_4O_4$ 530.55</p> <p>Line 7 of Coating the Solid Phase—Im</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
E-LINKED IMM UNOSORBENT ASSAY (ELISA)							<i>mobilization of Capture Reagent. Change 1–10 µg/well to: 1–10 µg/mL</i>
ESZOPICLONE ADDITIONAL REQUIREMENT S/USP Reference Standards <11>	USP40–NF35	4090	26-Jan-2018	1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	Line 2 of USP Eszopiclone Related Compound A RS: Change 6-(5-Chloropyridin-2-yl)-7-oxo-6,7-dihydro-5H-pyridolo[3,4-b]pyrazin-5-yl 4-methylpiperazine-1-carboxylate 4-oxide. C ₁₇ H ₁₇ ClN ₆ O ₄ 404.81 to: [Note—This material may be available in the free base or salt form.] 6-(5-Chloropyrid

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
POWDERED IDENTIFICATION OF ECHINACEA ANGUSTIFOLIA	USP40–NF35	6926	26-Jan-2018	1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	in-2-yl)-7-oxo-6,7-di hydro-5H -pyr rolo[3,4 -b]pyrazin-5-yl 4 -methylpiperazine-1-carboxylate 4-oxide. $C_{17}H_{17}ClN_6O_4$ 404.81 6-(5-Chloropyridin-2-yl)-7-oxo-6,7-di hydro-5H -pyr rolo[3,4 -b]pyrazin-5-yl 4 -methylpiperazine-1-carboxylate 4-oxide, 3-chlorobenzoic salt (1:1). $C_{17}H_{17}ClN_6O_4 \cdot C_7H_5ClO$ 561.38 Line 5 of A. Thin-Layer Chromatography/Presence of echinacoside and

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
							<p><i>dcaffeoylquinic acid/System suitability.</i></p> <p>Change <i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to:</p> <p><i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R</p>

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ECHINACEA PURPUREA AERIAL PARTS	IDENTIFICATIO N	USP40–NF35	6937	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	F) that are clearly separated, AND Line 3 of C.: Change <i>Standard solution B</i> , to: <i>Standard solution C</i> , Line 3 of A. <i>Thin-Layer Chromatography/ Presence of chicoric acid and absence of echinacoside/System suitability.</i> Change <i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
ECHINACEA IDENTIFICATIO SPECIES DRY N EXTRACT CAPSULES	USP41–NF36	4590	26-Jan-2018	1-Feb-2018	USP42–NF37	Second Supplement to USP41–NF36	due to caftaric acid (lower R_F) that are clearly separated, to: <i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated, Line 4 of A. HPTLC for Articles of Botanical Origin <203>/For Capsules containing <i>Echinacea angustifolia</i> Dry Extract/System suitability. Change

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
QUETIAPINE E IM XTENDED- RELEASE TABLETS	PUR ITIES/Organic Impurities	<i>Revision Bulletin (Official November 01, 2017)</i>	Online	26-Jan-2018		1-Feb-2018	<i>USP42–NF37</i>	<i>Second Supplement to USP41–NF36</i>	<p><i>Standard solution B shows two major blue bands at about the middle section due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to:</i></p> <p><i>Standard solution B shows two major blue bands at about the middle section due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated,</i></p> <p>Footnote a of Table 5: Change total impurities.</p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
DOBUTAMINE IN DEXTROSE INJECTION	<i>Identification</i> USP40–NF35	3843	26-Jan-2018	1-Feb-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	<p>to: total degradation products. AND Footnote b: Change total impurities. to: total degradation products. Line 1 of <i>B</i>: Change It meets the requirements for the <i>Identification</i> test under <i>Dextrose</i>. to: Add a few drops of a solution (1 in 20) to 5 mL of hot alkaline cupric tartrate TS. A copious red precipitate of cuprous oxide is formed.</p>
TIMOLOL	ADDITIONAL R USP40–NF35	6481	26-Jan-2018	1-Feb-2018	USP42–NF37	<i>Second</i>	Line 3 of USP

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
MALEATE TABLETS	EQUIREMENT S/USP Reference Standards <11>							<i>Supplement to USP41–NF36</i>	Timolol Related Compound D RS: Change C ₆ H ₉ N ₇ O ₄ S to: C ₆ H ₉ N ₃ O ₂ S
POWDERED ECHINACEA PALLIDA	IDENTIFICATIO N	USP40–NF35	6933	26-Jan-2018		1-Feb-2018	USP42–NF37	<i>Second Supplement to USP41–NF36</i>	Line 5 of A. <i>Thin-Layer Chromatography/Presence of echinacoside and absence of dicaffeoylquinic acid/System suitability.</i> Change Standard solution B shows two major blue bands at about the middle of the chromatogram due to caftaric acid (lower R _F) and chlorogenic acid (higher R _F) that are clearly separated, to:

Monograph Title	Section	Source Publication	Page Number	Errata Post Date	Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
POWDERED ECHINACEA	IDENTIFICATION	USP40–NF35	6944	26-Jan-2018		1-Feb-2018	USP42–NF37	Second Supplement to	<p><i>Standard solution B</i> shows two major blue bands at about the middle of the chromatogram due to caftaric acid (higher R_f) and chlorogenic acid (lower R_f) that are clearly separated, AND</p> <p>Lines 3 and 6 of C.: Change <i>Standard solution B</i>, to: <i>Standard solution C</i>, AND Change <i>Standard solution C</i>) to: <i>Standard solution D</i>)</p> <p>Line 3 of A. <i>Thin-Layer Chro</i></p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
PURPUREA EXTRACT						USP41–NF36	<p><i>matography/Pre sence of chicoric acid and absence of ec hinac oside/System suitability. Change Standard solution B shows two major blue bands at about the middle of the chromatogram due to caftaric acid (lower R_F) and chlorogenic acid (higher R_F) that are clearly separated, to: Standard solution B shows two major blue bands at about the middle of the</i></p>

Monograph Title Section	Source Publication	Page Number	Errata Post Date Sort ascending	Errata Official Date	Target Errata Print Publication	Target Online Fix Publication	Description
---	------------------------------------	-----------------------------	---	--------------------------------------	---	---	-------------

chromatogram due to caftaric acid (higher R_F) and chlorogenic acid (lower R_F) that are clearly separated,

Pagination

- [First page « First](#)
- [Previous page ‹ Previous](#)
- ...
- [Page 16](#)
- [Page 17](#)
- [Page 18](#)
- [Page 19](#)
- [Page 20](#)
- [Page 21](#)
- [Page 22](#)
- [Page 23](#)
- [Page 24](#)
- ...
- [Next page Next ›](#)
- [Last page Last »](#)