

Ganoderma Lucidum Fruiting Body

DEFINITION

Ganoderma Lucidum Fruiting Body consists of the dried fruiting body of *Ganoderma lucidum* (W. Curt.:Fr.) P. Karst. (Fam. Ganodermataceae). It contains NLT 0.3% of triterpenoid acids, calculated on the dried basis as a sum of ganoderic acids A, B, C₂, D, F, G, and H and ganoderenic acids B, C, and D.

IDENTIFICATION

Change to read:

• A. THIN-LAYER CHROMATOGRAPHY

Standard solution A: 1.0 mg/mL of USP Ganoderic Acid A RS in alcohol

Standard solution B: 0.3 mg/mL of USP Ergosterol RS in alcohol

Standard solution C: 50 mg/mL of USP Ganoderma Lucidum Fruiting Body Powdered Extract RS in alcohol. Sonicate for about 10 min, centrifuge, and use the supernatant.

Sample solution: Sonicate about 1 g of Ganoderma Lucidum Fruiting Body, finely powdered, in 50 mL of alcohol for 15 min. Centrifuge, withdraw the supernatant, and evaporate to dryness under reduced pressure at 50°. Dissolve the residue in 2.0 mL of alcohol, centrifuge, and use the supernatant.

Chromatographic system

(See *Chromatography* (621), *Thin-Layer Chromatography*.)

Mode: HPTLC

Adsorbent: Chromatographic silica gel with an average particle size of 5- μ m (HPTLC plate).¹ Predevelop the plate in methanol, and dry at 105° for 30 min.

Application volume: 2 μ L each of *Standard solution A* and *Standard solution B*, and 4 μ L each of *Standard solution C* and the *Sample solution* as 8-mm bands

Column temperature: Ambient, not to exceed 30°

Developing solvent system: Toluene, ethyl formate, and formic acid (5: 5: 0.2)

Developing distance: 6 cm

Derivatization (IRA 1-Mar-2015) **reagent:** A solution of 10% sulfuric acid in alcohol. [NOTE—Prepare fresh. Slowly and gradually add sulfuric acid to ice-cold alcohol, and mix well.]

• System suitability

Samples: *Standard solution A*, *Standard solution B*, and *Standard solution C*

Suitability requirements

Chromatographic pattern: Under long-wave UV light (365 nm), the chromatogram of *Standard solution C* displays, in the bottom third of the plate, the following bands in the order of increasing *R_f*: a yellowish or orange band (sometimes, two orange bands are seen); a bluish-green band corresponding to the light-blue ganoderic acid A band in *Standard solution A*; an intense yellow band corresponding to ganoderic acid B, ganoderic acid G, ganoderic acid H, and ganoderenic acid B; and a bluish-green band coincident with ganoderic acid D and ganoderenic acid D. In the middle third of the chromatogram, a variable number of blue-green bands appear. At the top of the middle third of the *Standard solution C* chromatogram, a somewhat diffuse band coincident with the ergosterol band in *Standard solution B* is

seen. In the upper third of the chromatogram, three or four diffuse bands of varying colors appear. Under white light, *Standard solution C* exhibits, in its lower third, two brownish-red bands, the upper of them coincident with the ganoderic acid A band in *Standard solution A*, followed by a more intense brown band; and a lighter brown band corresponding to ganoderic acid D and ganoderenic acid D. In the middle third of the chromatogram, five or six light-brown bands are seen; one of those, deepest in color and relatively diffuse, corresponds to the ergosterol band in *Standard solution B*. Two or three light-brown bands are seen under white light in the upper third of the chromatogram of *Standard solution C*.

[NOTE—The *Standard solutions* are stable for 72 h at room temperature.]

Analysis

Samples: *Standard solution A*, *Standard solution B*, *Standard solution C*, and *Sample solution*

Apply the samples as bands and dry in air. Develop in a saturated chamber, remove the plate, air-dry, treat with *Derivatization reagent*, and heat at 105°–110° for 5 min. Immediately examine under white light and under the long-wave UV light (365 nm).

Acceptance criteria: Under long-wave UV light (365 nm) and under white light, the chromatogram of the *Sample solution* exhibits bands corresponding in color and *R_f* to similar bands in the chromatogram of *Standard solution C*, at the *R_f* values listed for *System suitability*. Under white light, the chromatogram of the *Sample solution* exhibits an additional violet band above the ergosterol band. [NOTE—The *Sample solution* is stable for 72 h at room temperature.] (IRA 1-Mar-2015)

• B. HPLC

Analysis: Proceed as directed in the test for *Content of Triterpenoid Acids*.

Acceptance criteria: The chromatogram of the *Sample solution* exhibits peaks at the retention times corresponding to those of ganoderenic acid C, ganoderic acid C₂, ganoderic acid G, ganoderenic acid B, ganoderic acid B, ganoderic acid A, ganoderic acid H, ganoderenic acid D, ganoderic acid D, and ganoderic acid F in the chromatogram of *Standard solution B*.

• C. HPLC

Analysis: Proceed as directed in the test for *Content of Water-Soluble Polysaccharides*.

Acceptance criteria: The chromatogram of the *Sample solution* exhibits peaks at the retention times corresponding to the peaks due to mannose, glucuronic acid, dextrose, galactose, and L-fucose in the chromatogram of the *Standard solution*.

COMPOSITION

Change to read:

• CONTENT OF TRITERPENOIC ACIDS

Solution A: 0.075% phosphoric acid in water

Solution B: Acetonitrile

Mobile phase: See *Table 1*.

Table 1

Time (min)	Solution A (%)	Solution B (%)
0	80.0	20.0
3	73.5	26.5
34	73.5	26.5
52	61.5	38.5
53	80.0	20.0
58	80.0	20.0

¹ A suitable commercially available plate is the HPTLC Silica Gel 60 F₂₅₄ from EMD Millipore (e.g., Part No. 1.05642.0001).

2 Ganoderma

[NOTE—Maintain the *Mobile phase* at 73.5% of *Solution A* for the period sufficient for complete elution of ganoderic acid A.]

Standard solution A: 0.1 mg/mL of USP Ganoderic Acid A RS[●] (ERR 1-Dec-2014) in methanol. Sonicate to dissolve if necessary.

Standard solution B: Sonicate 40 mg of USP Ganoderma Lucidum Fruiting Body Powdered Extract RS in 5 mL of alcohol, and centrifuge. Pass through a nylon filter of 0.2- μ m pore size, and discard the initial 1 mL of the filtrate.

Sample solution: Transfer 2.0 g of Ganoderma Lucidum Fruiting Body, finely powdered and accurately weighed, to a 200-mL round-bottom flask, add 75 mL of alcohol, attach a condenser, reflux for 45 min, cool, and filter. Rinse the flask with two 10-mL portions of alcohol and filter, combining the rinsates and the filtrate. Evaporate to dryness under reduced pressure, and dissolve the residue in about 20 mL of alcohol. Transfer the solution to a 25-mL volumetric flask, dilute with alcohol to volume, and mix well. Pass through a nylon filter of 0.2- μ m pore size, and discard the initial 1 mL of the filtrate. [NOTE—To facilitate the chromatographic column longevity, the following solid phase extraction procedure may be employed. Condition the solid phase extraction column containing about 200 mg of L1 packing with 5 mL of methanol followed by 3 mL of water; do not allow the column to dry. Transfer 2.0 mL of Ganoderma Lucidum Fruiting Body solution in alcohol into a 20-mL volumetric flask, dilute with water to volume, and mix well. Apply the entire volume onto the column, and elute at the rate of approximately 1 drop/s, employing vacuum. Rinse the column with 3 mL of water, and discard the rinsate. Elute with 2.0 mL of methanol and collect the eluate into the 2.0-mL volumetric flask. Adjust with methanol to volume, and mix well.]

[NOTE—This method may result in coelution of ganoderic acid A and ganoderic acid K.]

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 257 nm

Column: 2.1-mm \times 15-cm; 1.8- μ m packing L1

Column temperature: 25°

Flow rate: 0.4 mL/min

Injection volume: 5 μ L

System suitability

Samples: *Standard solution A* and *Standard solution B*

Suitability requirements

Chromatographic similarity: The chromatogram of *Standard solution B* is similar to the reference chromatogram provided with the lot of USP Ganoderma Lucidum Fruiting Body Powdered Extract RS being used.

Resolution: NLT 1.0 between ganoderic acid A and ganoderic acid H peaks, *Standard solution B*

Tailing factor: NMT 2.0 for the ganoderic acid A peak, *Standard solution A*

Relative standard deviation: NMT 2.0% determined from the ganoderic acid A peak in replicate injections, *Standard solution A*

Analysis

Samples: *Standard solution A*, *Standard solution B*, and *Sample solution*

[NOTE—*Standard solution A*, *Standard solution B*, and the *Sample solution* are stable for 24 h at room temperature.]

Using the chromatograms of *Standard solution A*, *Standard solution B*, and the reference chromatogram pro-

vided with the lot of USP Ganoderma Lucidum Fruiting Body Powdered Extract RS being used, identify all specified ganoderic and ganoderenic acids in the *Sample solution* chromatogram. The approximate relative retention times, with respect to ganoderic acid A, are provided in *Table 2*.

Table 2

Analyte	Relative Retention Time	Relative Response Factor
Ganoderenic acid C	0.36	0.51
Ganoderic acid C ₂	0.42	1.05
Ganoderic acid G	0.56	1.18
Ganoderenic acid B	0.60	0.45
Ganoderic acid B	0.66	1.10
Ganoderic acid A	1.00	1.00
Ganoderic acid H	1.05	1.54
Ganoderenic acid D	1.25	0.51
Ganoderic acid D	1.33	1.08
Ganoderic acid F	1.54	1.45

Separately calculate the percentages of each triterpenic acid in the portion of Ganoderma Lucidum Fruiting Body taken:

$$\text{Result} = (r_U/r_S) \times C_S \times (V/W) \times F \times 100$$

r_U = peak area of the relevant analyte from the *Sample solution*

r_S = peak area of ganoderic acid A in *Standard solution A*[●] (ERR 1-Dec-2014)

C_S = concentration of USP Ganoderic Acid A RS in *Standard solution A* (mg/mL)

V = volume of the *Sample solution* (mL)

W = weight of Ganoderma Lucidum Fruiting Body taken to prepare the *Sample solution* (mg)

F = relative response factor, with respect to ganoderic acid A (see *Table 2*)[●] (ERR 1-Dec-2014)

Calculate the sum of the percentages of all specified triterpenic acids.

Acceptance criteria

Sum of triterpenic acids: NLT 0.3% on the dried basis

CONTAMINANTS

• ELEMENTAL IMPURITIES—PROCEDURES (233)

Acceptance criteria

Arsenic: NMT 2.0 μ g/g

Cadmium: NMT 1.0 μ g/g

Lead: NMT 5.0 μ g/g

Mercury: NMT 1.0 μ g/g

• **ARTICLES OF BOTANICAL ORIGIN, General Method for Pesticide Residues Analysis (561):** Meets the requirements

• **MICROBIAL ENUMERATION TESTS (2021):** The total aerobic bacterial count does not exceed 10⁵ cfu/g, and the bile-tolerant Gram-negative bacteria count does not exceed 10³ cfu/g.

• **ABSENCE OF SPECIFIED MICROORGANISMS (2022):** Meets the requirements of the tests for absence of *Salmonella* species and *Escherichia coli*

SPECIFIC TESTS

• CONTENT OF WATER-SOLUBLE POLYSACCHARIDES

Solution A: 0.05 M phosphate buffer, pH 6.0

Solution B: Acetonitrile

Mobile phase: See *Table 3*.

Table 3

Time (min)	Solution A (%)	Solution B (%)
0	84.0	16.0
30	82.5	17.5
55	81.0	19.0
60	81.0	19.0
61	84.0	16.0

Reagent: 0.1 M solution of 1-phenyl-3-methyl-5-pyrazolone in methanol

Internal standard solution: 0.5 mg/mL of D-lyxose in water

Standard stock solution: Composite solution containing 0.20 mg/mL each of USP Mannose RS, USP D-Glucuronic Acid RS, and USP Galactose RS; 2.0 mg/mL of USP Dextrose RS; and 0.10 mg/mL of USP L-Fucose RS in water

Standard solution: Combine 0.125 mL of *Standard stock solution* with 0.125 mL of *Internal standard solution*, 0.300 mL of 0.15 M sodium hydroxide solution, and 0.50 mL of *Reagent* in a capped reaction vial. Seal the vial, heat at 70° for 30 min, and cool to room temperature. Add to the vial 0.300 mL of 0.15 M hydrochloric acid and 0.65 mL of water, mix well, and pass through a nylon filter of 0.45-µm or finer pore size.

[NOTE—The amounts of individual analytes (A_s) in the 0.125-mL aliquot of the *Standard solution* submitted to derivatization are approximately 0.25 mg for dextrose and 0.025 mg for mannose, galactose, and D-glucuronic acid.]

Sample solution: Transfer 2.0 g of Ganoderma Lucidum Fruiting Body, finely powdered and accurately weighed, into a 200-mL round-bottom flask, add 60 mL of water, and allow to stand for 1 h. Attach a condenser, heat under reflux for 4 h, and filter immediately. Transfer the residue and the filter to the same 200-mL round-bottom flask. Add 60 mL of water, heat under reflux for 3 h, and filter immediately. Rinse the flask with three 5-mL portions of water, and filter. Combine the filtrates and the rinsates in a 250-mL beaker, and evaporate on the water bath to dryness. Dissolve the residue in 5 mL of water, add 75 mL of alcohol, mix well, allow to stand at 4° for 12 h, and centrifuge at 4000 rpm for 30 min. Discard the supernatant, and dry the precipitate on a water bath. Dissolve the residue in hot water and quantitatively transfer into a 10-mL volumetric flask. Cool to room temperature, dilute with water to volume, and mix well. Centrifuge at 4000 rpm for 10 min. Accurately transfer 0.250 mL of the supernatant into a reaction vial, and add about 0.25 mL of 4 M trifluoroacetic acid. Seal the vial, heat at 110° for 4 h, cool to room temperature, add 0.5 mL of methanol, and evaporate to dryness at 60° under vacuum. Repeat the addition of 0.5 mL of methanol and subsequent evaporation three times. Add to the residue 0.125 mL of water, 0.125 mL of the *Internal standard solution*, 0.300 mL of 0.15 M sodium hydroxide solution, and 0.50 mL of the *Reagent*. Seal the vial, heat at 70° for 30 min, and cool to room temperature. Add to the vial 0.300 mL of 0.15 M hydrochloric acid and 0.65 mL of water, mix well, and pass through a nylon filter of 0.45-µm or finer pore size.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 250 nm

Column: 4.6-mm × 25-cm; 5-µm packing L1

Column temperature: 35°

Flow rate: 1.0 mL/min

Injection volume: 10 µL

System suitability

Sample: *Standard solution*

Suitability requirements

Resolution: NLT 1.5 between the D-lyxose peak and the closest subsequent peak, and NLT 1.5 between the glucuronic acid peak and the closest preceding peak

Tailing factor: NMT 2.0 for the dextrose peak

Relative standard deviation: NMT 2.0% determined for the dextrose peak in replicate injections

Analysis

Samples: *Standard solution* and *Sample solution*

[NOTE—*Standard solution* and *Sample solution* are stable for 24 h at room temperature.]

Using the chromatograms of the *Standard solution* and the reference chromatogram provided with the lot of USP Ganoderma Lucidum Fruiting Body Powdered Extract RS being used, identify the individual derivatized monosaccharides at about the following relative retention times, with respect to dextrose: 0.48 for mannose, 0.58 for lyxose, 0.82 for D-glucuronic acid, 1.09 for galactose, and 1.35 for L-fucose.

Separately calculate the percentages of derivatized monosaccharides in the portion of Ganoderma Lucidum Fruiting Body taken:

$$\text{Result} = (R_U/R_S) \times A_S \times (F/W) \times 100$$

R_U = peak response ratio of the relevant analyte to the internal standard from the *Sample solution*

R_S = peak response ratio of the relevant analyte to the internal standard from the *Standard solution*

A_S = amount of the relevant analyte in the aliquot of the *Standard solution* subjected to derivatization (mg)

F = dilution factor to account for the sample aliquot submitted to derivatization (0.250 mL) relative to the volume of the *Sample solution* (10.0 mL), 40

W = weight of Ganoderma Lucidum Fruiting Body taken to prepare the *Sample solution* (mg)

Calculate the sum of the percentages of mannose, D-glucuronic acid, dextrose, galactose, and L-fucose.

Acceptance criteria

Sum of monosaccharides: NLT 0.7% on the dried basis

Change to read:

• **BOTANICAL CHARACTERISTICS**

Macroscopic: Basidiocarp (fruiting body) morphology is highly variable. Shape of pileus (cap) ranges from reniform to subcircular, convex or concave, 1.5 cm or more broad, single to multiple layers thick (up to 3 cm); margin generally thick and blunt, sometimes acute. Pileus surface radially rugose (wrinkled) and concentrically sulcate; (ERR 1-Dec-2014) shiny, yellowish-red to reddish-black. Stipe (stem) attachment predominantly lateral; stipe length varies from very short to 10–12 cm long, 1–3 cm thick, cylindrical, reddish to almost black, laccate (lacquered). Hymenophore (pore surface) yellowish-white to tawny. Pores small, circular to irregular, 4–7 per mm, 6–200 µm in diameter, distance between axes of pores about 260 µm.

4 Ganoderma

Microscopic: Hyphal system trimitic with hyaline, thin-walled, clamped, septate generative hyphae, 1–4 μm in diameter, septa restricted to clamps, scantily branched, abundant at the growth margin of pileus and dissepiments (partitions). Skeletal hyphae are arboriform, aseptate, clampless, very long, 3–6 μm in diameter, scantily branched, branches with limited growth at distal end, with thick walls; they compose most of the context (flesh) and dissepiments, originating immediately behind the growth margin from generative hyphae. Binding hyphae of the “Bovista” type are aseptate, clampless, profusely branched, generally thinner and lighter than the skeletal, 1–3 μm in diameter. Basidiospores ovoid, double-walled, truncated at apex. Epispore thin, ovoid, hyaline, 9.0–11.5 \times 6.0–8.0 μm ; endospore thick, ovoid, 6.5–8.5 \times 5.0–6.5 μm , bearing relatively few long and thick echinules that support the epispore, sometimes fused into a short crest.

- **ARTICLES OF BOTANICAL ORIGIN, Foreign Organic Matter (561):** NMT 2.0%
- **LOSS ON DRYING (731)**
Sample: 1.0 g of powdered Ganoderma Lucidum Fruiting Body
Analysis: Dry at 105° for 4 h.
Acceptance criteria: NMT 17.0%
- **ARTICLES OF BOTANICAL ORIGIN, Total Ash (561)**
Sample: 1.0 g of powdered Ganoderma Lucidum Fruiting Body
Acceptance criteria: NMT 4.0%
- **ARTICLES OF BOTANICAL ORIGIN, Alcohol-Soluble Extractives, Method 1 (561)**

Sample: 2–4 g of powdered Ganoderma Lucidum Fruiting Body

Acceptance criteria: NLT 2.0%

- **ARTICLES OF BOTANICAL ORIGIN, Water-Soluble Extractives, Method 1 (561)**

Sample: 2–4 g of powdered Ganoderma Lucidum Fruiting Body

Acceptance criteria: NLT 3.0%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers, protected from light and moisture, and store at room temperature.
- **LABELING:** The label states the Latin binominal and, following the official name, the part of the fungus from which the article was derived.
- **USP REFERENCE STANDARDS (11)**
 - USP Dextrose RS
 - USP Ergosterol RS
 - USP L-Fucose RS
 - USP Galactose RS
 - USP Ganoderic Acid A RS
 - USP Ganoderma Lucidum Fruiting Body Powdered Extract RS
 - USP D-Glucuronic Acid RS
 - USP Mannose RS