

Estradiol Transdermal System

Type of Posting	Revision Bulletin
Posting Date	27–Jan–2017
Official Date	01–Feb–2017
Expert Committee	Chemical Medicines Monographs 5
Reason for Revision	Compliance

In accordance with the Rules and Procedures of the 2015-2020 Council of Experts, the Chemical Medicines Monographs 5 Expert Committee has revised the Estradiol Transdermal System monograph. The purpose for the revision is to add *Drug Release Test 4* to accommodate a drug product that was approved with different drug release test conditions and acceptance criteria.

The liquid chromatographic procedure used in *Drug Release Test 4* is based on analyses performed with the Symmetry C18 brand of L1 column manufactured by Waters. The typical retention time for estradiol is about 4.5 minutes.

The Estradiol Transdermal System Revision Bulletin supersedes the currently official monograph. The Revision Bulletin will be incorporated in the *Second Supplement to USP 40–NF 35*.

Should you have any questions, please contact Gerald Hsu, Ph.D., Senior Scientific Liaison (240-221-3097 or gdh@usp.org).

Estradiol Transdermal System

DEFINITION

Estradiol Transdermal System contains NLT 85.0% and NMT 120.0% of the labeled amount of estradiol ($C_{18}H_{24}O_2$).

IDENTIFICATION

- **A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

ASSAY

PROCEDURE

Diluent: Acetonitrile and water (1:1)

Mobile phase: Acetonitrile and water (55:45)

Standard solution: 0.1 mg/mL of USP Estradiol RS in *Diluent*

Sample solutions: Equivalent to 0.1 mg/mL of estradiol in *Diluent*, prepared as follows. Cut 10 Transdermal Systems into pieces, and keep the pieces from each system separate. Remove and discard the protective liners, if present, from the strips. Transfer the pieces of each system into separate stoppered flasks of suitable size, and add a measured volume of *Diluent* to each flask to provide the target estradiol concentration. Shake by mechanical means for about 3 h, and sonicate for 15 min.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 280 nm

Column: 4.6-mm \times 15-cm; packing L1

Column temperature: 35°

Flow rate: 1 mL/min

Injection size: 25 μ L

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: 0.9–1.6

Relative standard deviation: NMT 2.5%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of estradiol ($C_{18}H_{24}O_2$) in each Transdermal System taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Estradiol RS in the *Standard solution* (mg/mL)

C_U = nominal concentration of estradiol in the *Sample solution* (mg/mL)

Use the individual assays to determine *Uniformity of Dosage Units*.

Acceptance criteria: 85.0%–120.0%

OTHER COMPONENTS

ALCOHOL CONTENT (if present)

Diluent: Acetonitrile and water (1:1)

Internal standard solution: Prepare by diluting 4.0 mL of dehydrated methanol with water to 100 mL.

Standard stock solution: 5.0 mg/mL of ethanol in *Diluent*. Prepare by weighing by difference 1.6 mL of dehydrated alcohol into a tared 50-mL volumetric flask containing 15 mL of water, and dilute with *Diluent* to volume. Pipet 10.0 mL of this solution into a 50-mL volumetric flask, and dilute with *Diluent* to volume.

Standard solution: 2.5 mg/mL of ethanol. Prepare by pipeting 25.0 mL of the *Standard stock solution* into a 50-mL volumetric flask. Add 5.0 mL of the *Internal standard solution*, and dilute with water to volume.

Sample solutions: Prepare as directed for the *Sample solutions* in the *Assay*, with the following changes. Pipet 25.0 mL of each solution into individual 50-mL volumetric flasks. Add 5.0 mL of the *Internal standard solution*, and dilute with water to volume.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: GC

Detector: Flame ionization

Column: 2-mm \times 2-m glass; support S2

Temperature

Column: 100°

Injection port: 200°

Detector: 200°

Carrier gas: Helium

Flow rate: 30 mL/min

Injection size: 2 μ L

System suitability

Sample: *Standard solution*

[NOTE—The relative retention times for the methanol and alcohol peaks are 0.4 and 1.0, respectively.]

Suitability requirements

Relative standard deviation: NMT 1.5% from the peak response ratio of alcohol to methanol

Analysis

Samples: *Standard solution* and *Sample solutions*

Calculate the percentage of alcohol (C_2H_5OH) in each Transdermal System taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

R_U = peak response ratio of alcohol to methanol from the *Sample solution*

R_S = peak response ratio of alcohol to methanol from the *Standard solution*

C_S = concentration of dehydrated alcohol in the *Standard solution* (mg/mL)

C_U = nominal concentration of alcohol in the *Sample solution* (mg/mL)

Average the percentage of alcohol found in the Transdermal Systems analyzed.

Acceptance criteria: 80%–120% of the labeled amount of C_2H_5OH

PERFORMANCE TESTS

Change to read:

DRUG RELEASE (724)

Test 1: For products labeled for dosing every 84 h

Medium: Water; 900 mL, deaerated

Apparatus 5: 50 rpm

Times: 24, 48, and 96 h

Mobile phase: Water and acetonitrile (3:2)

Standard solution: 9 μ g/mL of USP Estradiol RS in dehydrated alcohol. Dilute this solution with *Medium* to obtain solutions having concentrations of about 0.9, 0.45, and 0.045 μ g/mL.

Sample solution: At each sampling time interval, withdraw a 10-mL aliquot of the solution under test.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: Fluorimetric, with excitation at 220 nm and emission at 270 nm

2 Estradiol

Column: 4.6-mm × 3-cm; packing L1
Temperature: 40°
Flow rate: 1.0 mL/min
Injection size: 50 µL
System suitability
Sample: *Standard solution*
Tailing factor: 0.9–2.5
Relative standard deviation: NMT 3.0%, using 0.45 µg/mL of the *Standard solution*

Analysis: Plot the peak responses of the *Standard solutions* versus concentration, in µg/mL, of estradiol. From the graph determine the amount, in µg/mL, of estradiol released. Calculate the cumulative release rate as percentage of the labeled amount of estradiol: At 24 h:

$$\text{Result} = \{[900(A_1 - b)] / (1000 \times m \times L)\} \times 100$$

At 48 h:

$$\text{Result} = \{[890(A_2 - b) + 10(A_1 - b)] / (1000 \times m \times L)\} \times 100$$

At 96 h:

$$\text{Result} = \{[880(A_3 - b) + 10(A_2 - b) + 10(A_1 - b)] / (1000 \times m \times L)\} \times 100$$

A_1 = peak area of estradiol in the *Sample solution* at the first time interval

A_n = peak area of estradiol in the *Sample solution* at the release interval n

m = slope of the calibration curve

b = y-intercept of the calibration curve

L = Transdermal System label claim (mg)

Tolerances: See Table 1.● (RB 1-Feb-2017)

Table 1

Time (h)	Amount Dissolved (release rate)
24	1.2%–6.0%
48	3.0%–11.4%
96	5.0%–16.3%

●The percentages of the labeled amount of estradiol ($C_{18}H_{24}O_2$) released at the times specified conform to *Drug Release (724)*, *Acceptance Table 1.●* (RB 1-Feb-2017)

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Drug Release Test 2*.

Medium: 0.005 M phosphate buffer, pH 5.5, containing 0.3% sodium lauryl sulfate; 500 mL

Apparatus 5: 100 rpm. Use a 76-mm stainless steel disk assembly. Adhere the patch to the disk assembly using transfer tape. [NOTE—A suitable tape is available as 3M adhesive transfer tape 927, www.mmm.com.]

Times: 1, 4, 8, and 24 h

Mobile phase: Acetonitrile and water (1:1)

Standard stock solution: 800 µg/mL of USP Estradiol RS in acetone

Standard solution: Dilute the *Standard stock solution* with *Medium* to obtain a solution having a known concentration close to that expected in the solution under test, assuming 100% drug release.

Sample solution: At each sampling time interval, withdraw a known volume aliquot of the solution under test.

Chromatographic system

(See *Chromatography (621)*, *System Suitability*.)

Mode: LC
Detector: UV 205 nm
Column: 3.9-mm × 30-cm; packing L1
Flow rate: 1.0 mL/min
Injection size: 100 µL

System suitability

Sample: *Standard solution*

Tailing factor: NMT 2.0

Relative standard deviation: NMT 3.0%

Analysis: Calculate the amount of estradiol released at each sampling time:

$$M_i = (r_u/r_s) \times C_s \times V_i$$

$$m_1 = M_1$$

$$m_2 = M_2 + M_1(V_a/V_1)$$

$$m_3 = M_3 + M_2(V_a/V_2) + M_1(V_a/V_1)$$

$$m_4 = M_4 + M_3(V_a/V_3) + M_2(V_a/V_2) + M_1(V_a/V_1)$$

Calculate the percentage of the labeled amount of estradiol released at each sampling time:

$$\text{Result} = (m_i/L) \times 100$$

M_i = amount of estradiol released into the *Medium* at a given sampling time (mg)

r_u = peak response from the *Sample solution*

r_s = peak response from the *Standard solution*

C_s = concentration of the *Standard solution* (mg/mL)

V_i = corrected volume of the *Medium* at a given sampling time (mL)

m_1, m_2, m_3, m_4 = total amounts of estradiol released from the patch at given sampling times (mg)

M_1, M_2, M_3, M_4 = amounts of estradiol released into the *Medium* at given sampling times (mg)

V_a = volume of the aliquot taken from the dissolution vessel at each sampling time (mL)

V_1, V_2, V_3 = volumes of *Medium* at given sampling times (mL)

L = Transdermal System label claim (mg)

Tolerances: See Table 2.● (RB 1-Feb-2017)

Table 2

Time (h)	Amount Dissolved (release rate)
1	15%–40%
4	45%–70%
8	70%–90%
24	NLT 80%

●The percentages of the labeled amount of estradiol ($C_{18}H_{24}O_2$) released at the times specified conform to *Drug Release (724)*, *Acceptance Table 1.●* (RB 1-Feb-2017)

Test 3: If the product complies with this test, the labeling indicates that it meets USP *Drug Release Test 3*.

Medium: 1% (v/v) polysorbate 40 in water; 900 mL

Apparatus 5: 50 rpm

Times: 4, 8, and 24 h

Standard stock solution: Known concentration (mg/mL) of USP Estradiol RS in methanol

Standard solution: Five different concentrations within the range of the expected release amounts of estradiol, prepared as follows. Add 1.0 mL of polysorbate 40 into a 100-mL volumetric flask, and then add the required amount of *Standard stock solution*. Mix well to dissolve the polysorbate 40, and dilute with water to volume.

Sample solution: At each sampling time interval, withdraw a known volume aliquot of the solution under test.

Mobile phase: Acetonitrile and water (2:3)

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 225 nm

Column: 4.6-mm × 15-cm, 5-μm packing L1 for 9-cm² systems; 4.6-mm × 12.5-cm, 5-μm packing L1 for 18-, 27-, or 36-cm² systems. In any case, a guard column containing packing L1 is used.

Flow rate: 1.0 mL/min

Injection size: 50 μL

System suitability

Sample: *Standard solution*

Relative standard deviation: NMT 2.0%

Analysis: Calculate the cumulative release rate as a percentage of the labeled amount of estradiol:

$$\text{Result} = \{[900(A - b)] / (1000 \times m \times L)\} \times 100$$

A = peak area of estradiol in the *Sample solution* at each time interval

b = y-intercept of the calibration curve

m = slope of the calibration curve

L = Transdermal System label claim (mg)

Tolerances: The percentages of the labeled amount of estradiol (C₁₈H₂₄O₂) released at the times specified conform to *Table 3*, *Table 4*, and *Table 5*. (RB 1-Feb-2017) L1 (6 units)

Table 3

Time (h)	Amount Dissolved (individual values)
4	40%–71%
8	58%–94%
24	NLT 75%

L2 (12 units)

Table 4

Time (h)	Amount Dissolved (average of 12)	Amount Dissolved (individual values)
4	40%–71%	34%–77%
8	58%–94%	50%–102%
24	NLT 75%	NLT 68%

L3 (24 units)

Table 5

Time (h)	Amount Dissolved (average of 24)	Amount Dissolved (individual for 22 units of 24)	Amount Dissolved (individual for 24)
4	40%–71%	34%–77%	29%–82%
8	58%–94%	50%–102%	43%–109%
24	NLT 75%	NLT 68%	NLT 60%

Test 4: If the product complies with this test, the labeling indicates that it meets USP *Drug Release Test 4*.

Medium: Water; 500 mL for 0.025 mg/day and 0.0375 mg/day dosage; 900 mL for 0.05 mg/day, 0.075 mg/day, and 0.1 mg/day dosage

Apparatus 6: 50 rpm. Use a stainless steel cylinder assembly. Adhere the Transdermal System to the cylinder assembly using a strip of suitable double-sided transfer tape.

Times: 2, 6, and 12 h

Buffer solution: 25 mM of monobasic sodium phosphate, adjusted with phosphoric acid to a pH of 3.0

Mobile phase: Acetonitrile and *Buffer solution* (40:60)

Standard stock solution: 0.2 mg/mL of USP Estradiol RS in methanol

Standard solution: Dilute the *Standard stock solution* with *Medium* to obtain a solution having a known concentration that is approximately 90% of the concentration expected from complete release in the solution under test.

Sample solution: At each sampling time interval, withdraw about 1.5 mL of the solution under test. Place each sample aliquot into an amber HPLC vial.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 280 nm

Column: 3.0-mm × 10-cm; 3.5-μm packing L1

Flow rate: 0.5 mL/min

Injection volume: 15 μL

Run time: 2.5 times the retention time of estradiol

System suitability

Sample: *Standard solution*

Tailing factor: NMT 1.8

Relative standard deviation: NMT 3.0%

Analysis

Samples: *Standard solution* and *Sample solution*
 Calculate the concentration (*C_i*) of estradiol (C₁₈H₂₄O₂) in the sample withdrawn from the vessel at time point *i*:

$$C_i = (r_i/r_s) \times C_s$$

r_i = peak response of estradiol from the *Sample solution* at time point *i*

r_s = peak response of estradiol from the *Standard solution*

C_s = concentration of USP Estradiol RS in the *Standard solution* (mg/mL)

Calculate the percentage of the labeled amount of estradiol (C₁₈H₂₄O₂) released at each time point (*i*):

$$\text{Result}_1 = C_i \times V \times (1/L) \times 100$$

$$\text{Result}_2 = \{[C_2 \times (V - V_s)] + (C_1 \times V_s)\} \times (1/L) \times 100$$

$$\text{Result}_3 = \{[C_3 \times [V - (2 \times V_s)]] + [(C_2 + C_1) \times V_s]\} \times (1/L) \times 100$$

C_i = concentration of estradiol in the portion of the sample withdrawn at each time point (*i*) (mg/mL)

V = volume of *Medium*, 900 or 500 mL

L = Transdermal System label claim (mg)

V_s = volume of *Sample solution* withdrawn from the *Medium* (mL)

Tolerances: See *Table 6*.

4 Estradiol

Table 6

Time (h)	Amount Dissolved (release rate, %)
2	20–40
6	48–68
12	70–90

The percentages of the labeled amount of estradiol ($C_{18}H_{24}O_2$) released at the times specified conform to *Dissolution* (711), *Acceptance Table 2*. (RB 1-Feb-2017)

- **UNIFORMITY OF DOSAGE UNITS (905):** The results from the Transdermal Systems used in the *Assay* meet the requirements.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in hermetic, light-resistant, unit-dose pouches.

Change to read:

- **LABELING:** The label states the total amount of estradiol in the Transdermal System and the release rate, in mg/day, for the duration of application of one system. When more than one *Drug Release* test is given, the labeling states the *Drug Release* test used only if *Test 1* is not used.

(RB 1-Feb-2017)

- **USP REFERENCE STANDARDS (11)**
 USP Estradiol RS