

Potassium Chloride Extended-Release Tablets

| | |
|----------------------------|---------------------------------|
| Type of Posting | Revision Bulletin |
| Posting Date | 31-Jul-2019 |
| Official Date | 01-Aug-2019 |
| Expert Committee | Chemical Medicines Monographs 5 |
| Reason for Revision | Compliance |

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 5 Expert Committee has revised the Potassium Chloride Extended-Release Tablets monograph. The purpose for the revision is to add *Dissolution Test 4* to accommodate FDA-approved drug products with different dissolution conditions and tolerances than the existing dissolution tests.

The Potassium Chloride Extended-Release Tablets Revision Bulletin supersedes the currently official monograph.

The addition of *Dissolution Test 3* to the Potassium Chloride Extended-Release Tablets monograph is currently being proposed under the Pending Monograph process.

Should you have any questions, please contact Ren-Hwa Yeh, Senior Scientific Liaison (301-998-6818 or rhy@usp.org).

Potassium Chloride Extended-Release Tablets

DEFINITION

Potassium Chloride Extended-Release Tablets contain NLT 90.0% and NMT 110.0% of the labeled amount of potassium chloride (KCl).

IDENTIFICATION

- **A. IDENTIFICATION TESTS—GENERAL** (191), *Chemical Identification Tests, Potassium*

Sample solution: A portion of the filtrate, obtained as directed for the designated *Sample stock solution* in the Assay

Acceptance criteria: Meet the requirements

- **B. IDENTIFICATION TESTS—GENERAL** (191), *Chemical Identification Tests, Chloride*

Sample solution: A portion of the filtrate, obtained as directed for the designated *Sample stock solution* in the Assay

Acceptance criteria: Meet the requirements

ASSAY

PROCEDURE

[NOTE—If necessary, first score nonsugar-coated Tablets. Retain a portion of the filtrate of either *Sample stock solution 1* or *Sample stock solution 2* for use in *Identification A* and *B*.]

Standard stock solution: 19.07 µg/mL of potassium chloride, previously dried at 105° for 2 h, in water. This solution contains 10 µg/mL of potassium.

Standard solutions: To separate 100-mL volumetric flasks transfer 10.0, 15.0, and 20.0 mL, respectively, of *Standard stock solution*. To each flask add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume. The *Standard solutions* contain 1.0, 1.5, and 2.0 µg/mL of potassium, respectively.

Sample preparation 1

Sample stock solution 1: Nominally 0.06 mg/mL of potassium chloride prepared as follows. Place NLT 20 Tablets in a suitable container with 400 mL of water, heat to boiling, and boil for 20 min. Allow to cool, transfer the solution to a 1000-mL volumetric flask, and dilute with water to volume. Filter and discard the first 20 mL of the filtrate. Transfer a measured volume of the subsequent filtrate, equivalent to 60 mg of potassium chloride, to a 1000-mL volumetric flask, and dilute with water to volume.

Sample solution 1: Nominally 3 µg/mL of potassium chloride prepared as follows. Transfer 5.0 mL of *Sample stock solution 1* to a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

Sample preparation 2 (for formulations containing crystals coated with hydrophobic polymers)

Sample stock solution 2: Nominally 0.06 mg/mL of potassium chloride prepared as follows. Place NLT 20 Tablets in a 2000-mL volumetric flask. Add 1200 mL of a mixture of acetonitrile and water (1:1), and shake by mechanical means, or stir using a magnetic bar for 90 min. Dilute with the mixture of acetonitrile and water (1:1) to volume. Allow to stand for 90 min. Pass through a filter of 0.2-µm pore size. Transfer a measured volume of the filtrate, and quantitatively dilute with water to obtain a solution with a concentration of 0.06 mg/mL. [NOTE—Alternatively, *Sample stock solution 2* can be prepared by the following procedure. Nominally 0.15 mg/mL of potassium chloride from NLT 20 finely

powdered Tablets, prepared as follows. Transfer an appropriate amount of the powder, equivalent to about 5–6 Tablets, to a suitable volumetric flask, add 10% of the final flask volume of acetone, and sonicate for 45 min with intermittent shaking. Add 80% of the final flask volume of water and sonicate for 45 min with intermittent shaking. Cool to room temperature and dilute with water to volume. Centrifuge a portion of the solution at 5000 rpm for 10 min. Transfer an appropriate amount of the supernatant to a 100-mL volumetric flask and dilute with water to volume to obtain a solution with a concentration of 0.15 mg/mL.]

Sample solution 2: Nominally 3 µg/mL of potassium chloride prepared as follows. Transfer an appropriate amount of *Sample stock solution 2* to a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

Instrumental conditions

(See *Atomic Absorption Spectroscopy* (852).)

Mode: Atomic absorption spectrophotometry

Analytical wavelength: Potassium emission line at 766.5 nm

Lamp: Potassium hollow-cathode

Flame: Air–acetylene

Blank: Water

Analysis

Samples: *Standard solutions*, *Sample solution 1* or *Sample solution 2*, and *Blank*

Plot the absorbances of the *Standard solutions* versus the concentration of potassium, in µg/mL, and draw the straight line best fitting the three plotted points. From the graph, determine the concentration of potassium in the *Sample solution* (µg/mL).

Calculate the percentage of the labeled amount of potassium chloride (KCl) in each Tablet taken:

$$\text{Result} = (C/C_U) \times (M_r/A_r) \times 100$$

- C = concentration of potassium in the *Sample solution* as determined in this test (µg/mL)
C_U = nominal concentration of potassium chloride in the *Sample solution* (µg/mL)
M_r = molecular weight of potassium chloride, 74.55
A_r = atomic weight of potassium, 39.10

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Change to read:

DISSOLUTION (711)

Test 1

Medium: Water; 900 mL

Apparatus 2: 50 rpm

Time: 2 h

Standard stock solution: 19.07 µg/mL of potassium chloride, previously dried at 105° for 2 h, in water. This solution contains 10 µg/mL of potassium.

Standard solutions: To separate 100-mL volumetric flasks transfer 10.0, 15.0, and 20.0 mL, respectively, of *Standard stock solution*. To each flask add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume. The *Standard solutions* contain, respectively, 1.0, 1.5, and 2.0 µg/mL of potassium.

Sample stock solution: Filter the solution under test, and dilute with *Medium* to obtain a solution containing nominally 60 µg/mL of potassium chloride.

Sample solution: Transfer 5.0 mL of the *Sample stock solution* to a 100-mL volumetric flask. Add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

Instrumental conditions

(See *Atomic Absorption Spectroscopy* (852).)

Mode: Atomic absorption spectrophotometry

Analytical wavelength: Potassium emission line at 766.5 nm

Lamp: Potassium hollow-cathode

Flame: Air-acetylene

Blank: Water

Analysis

Samples: *Standard solutions*, *Sample solution*, and *Blank*
Plot the absorbances of the *Standard solutions* versus the concentration of potassium, in µg/mL, and draw the straight line best fitting the three plotted points. From the graph, determine the concentration of potassium in the *Sample solution* (µg/mL).

Calculate the percentage of the labeled amount of potassium chloride (KCl) dissolved:

$$\text{Result} = [C \times D \times (V/L)] \times (M_r/A_r) \times 100$$

C = concentration of potassium in the *Sample solution* as determined in this test (µg/mL)

D = dilution factor of the *Sample solution*

V = volume of *Medium*, 900 mL

L = labeled amount of potassium chloride (µg/ Tablet)

M_r = molecular weight of potassium chloride, 74.55

A_r = atomic weight of potassium, 39.10

Tolerances: NMT 35% (*Q*) of the labeled amount of potassium chloride (KCl) is dissolved in 2 h. The requirements are met if the quantities dissolved from the Tablets tested conform to *Table 1* instead of the table shown in *Dissolution* (711).

Table 1

| Stage | Number Tested | Acceptance Criteria |
|-----------------------|---------------|--|
| <i>S</i> ₁ | 6 | Each unit is within the range $Q \pm 30\%$. |
| <i>S</i> ₂ | 6 | Average of 12 units (<i>S</i> ₁ + <i>S</i> ₂) is within the range between $Q - 30\%$ and $Q + 35\%$, and no unit is outside the range $Q \pm 40\%$. |
| <i>S</i> ₃ | 12 | Average of 24 units (<i>S</i> ₁ + <i>S</i> ₂ + <i>S</i> ₃) is within the range between $Q - 30\%$ and $Q + 35\%$, and NMT 2 units are outside the range $Q \pm 40\%$. |

Test 2: If the product complies with this procedure, the labeling indicates that it meets USP *Dissolution Test 2*.

Standard stock solution and Standard solutions:

Prepare as directed in *Test 1*.

Medium: Water; 900 mL

Apparatus 2: 50 rpm

Times: 1, 2, 4, and 8 h

Sample stock solution: Transfer 4.0 mL of the solution under test into either a 50-mL volumetric flask (for 750-mg Tablet) or a 100-mL volumetric flask (for 1500-mg Tablet), dilute with water to volume, and filter.

Sample solution: Transfer 4.0 mL of the *Sample stock solution* to a 100-mL volumetric flask. Add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

Blank solution: To a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

Instrumental conditions: Proceed as directed in *Test 1*, except do not use the *Blank*.

System suitability

Samples: *Standard solutions*

Suitability requirements

Linearity: Correlation coefficient NLT 0.99

Relative standard deviation: NMT 5.0% from 5 replicate analyses of the 1.5-µg/mL *Standard solution*

Analysis

Samples: 1.5-µg/mL *Standard solution*, *Sample solution*, and *Blank solution*

Calculate the percentage of the labeled amount of potassium chloride (KCl) dissolved:

$$\text{Result}_i = [(A_U/A_S) \times C_S \times D \times (V/L)] \times (M_r/A_r) \times 100$$

A_U = absorbance of potassium in the *Sample solution*

A_S = absorbance of potassium in the *Standard solution*

C_S = concentration of potassium in the *Standard solution* (µg/mL)

D = dilution factor of the *Sample solution*

V = volume of *Medium*, 900 mL

L = labeled amount of potassium chloride (µg/ Tablet)

M_r = molecular weight of potassium chloride, 74.55

A_r = atomic weight of potassium, 39.10

Tolerances: See *Table 2*.

Table 2

| Time Point (<i>t</i>) | Time (h) | Amount Dissolved (%) | |
|-------------------------|----------|----------------------|----------------|
| | | 750 mg/Tablet | 1500 mg/Tablet |
| 1 | 1 | 10–30 | 5–25 |
| 2 | 2 | 30–50 | 25–45 |
| 3 | 4 | 60–80 | 55–75 |
| 4 | 8 | NLT 80 | NLT 85 |

The percentages of the labeled amount of potassium chloride (KCl), dissolved at the times specified, conform to *Dissolution* (711), *Acceptance Table 2*.

▲Test 4: If the product complies with this procedure, the labeling indicates that it meets USP *Dissolution Test 4*.

Standard stock solution and Instrumental conditions:

Proceed as directed in *Test 1*, except *Blank*.

Medium: Water; 900 mL, degassed

Apparatus 2: 50 rpm

Times: 2, 4, and 8 h

Sodium chloride solution: 0.2 g/mL of sodium chloride in water

Hydrochloric acid solution: Dilute 100 mL of hydrochloric acid with 300 mL of water.

Standard solutions: To separate 100-mL volumetric flasks transfer 10.0, 15.0, and 20.0 mL, respectively, of *Standard stock solution*. To each flask add 2.0 mL of *Sodium chloride solution* and 4.0 mL of *Hydrochloric acid solution*, and dilute with water to volume. The *Standard*

solutions contain 1.0, 1.5, and 2.0 µg/mL of potassium, respectively.

Sample stock solution: Pass a portion of the solution under test through a filter with a suitable pore size and use the filtrate.

Sample solution: Transfer 1.0 mL of the *Sample stock solution* to a suitable volumetric flask and dilute with water if necessary. To the final dilution, add 2.0% flask volume of *Sodium chloride solution* and 4.0% flask volume of *Hydrochloric acid solution*, and dilute with water to volume.

Blank: To a suitable volumetric flask, add 2.0% flask volume of *Sodium chloride solution* and 4.0% flask volume of *Hydrochloric acid solution*, and dilute with water to volume.

System suitability

Samples: *Standard solutions*

Suitability requirements

Linearity: Correlation coefficient NLT 0.999

Relative standard deviation: NMT 1.5% from the absorbance responses of 5 replicate analyses of each *Standard solution*

Analysis: Proceed as directed in *Test 1*.

Tolerances: See *Table 3*.

Table 3

| Time Point (l) | Time (h) | Amount Dissolved (%) |
|----------------|----------|----------------------|
| 1 | 2 | 22–42 |
| 2 | 4 | 44–64 |
| 3 | 8 | NLT 80 |

The percentages of the labeled amount of potassium chloride (KCl), dissolved at the times specified, conform to *Dissolution (711)*, *Acceptance Table 2*.▲ (RB 1-Aug-2019)

- **UNIFORMITY OF DOSAGE UNITS (905):** Meet the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, and store at a temperature not exceeding 30°.
- **LABELING:** The label states with which *Sample preparation* in the *Assay* the product complies only if *Sample preparation 1* is not used. When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.